



An inverse problem for the magnetic Schrödinger equation in infinite cylindrical domains

M Bellassoued, Y Kian, Eric Soccorsi

► To cite this version:

M Bellassoued, Y Kian, Eric Soccorsi. An inverse problem for the magnetic Schrödinger equation in infinite cylindrical domains. Publications of the Research Institute for Mathematical Sciences, 2018, 54 (4), pp.679-728. 10.4171/PRIMS/54-4-1 . hal-01319127

HAL Id: hal-01319127

<https://hal.science/hal-01319127>

Submitted on 20 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AN INVERSE PROBLEM FOR THE MAGNETIC SCHRÖDINGER EQUATION IN INFINITE CYLINDRICAL DOMAINS

M. BELLASSOUED, Y. KIAN, AND E. SOCCORSI

ABSTRACT. We study the inverse problem of determining the magnetic field and the electric potential entering the Schrödinger equation in an infinite 3D cylindrical domain, by Dirichlet-to-Neumann map. The cylindrical domain we consider is a closed waveguide in the sense that the cross section is a bounded domain of the plane. We prove that the knowledge of the Dirichlet-to-Neumann map determines uniquely, and even Hölder-stably, the magnetic field induced by the magnetic potential and the electric potential. Moreover, if the maximal strength of both the magnetic field and the electric potential, is attained in a fixed bounded subset of the domain, we extend the above results by taking finitely extended boundary observations of the solution, only.

Keywords: Inverse problem, magnetic Schrödinger equation, Dirichlet-to-Neumann map, infinite cylindrical domain.

1. INTRODUCTION

1.1. Statement of the problem. Let ω be a bounded connected domain of \mathbb{R}^2 with C^2 boundary $\partial\omega$, and set $\Omega := \omega \times \mathbb{R}$. For $T > 0$, we consider the initial boundary value problem (IBVP)

$$\begin{cases} (i\partial_t + \Delta_A + q)u = 0, & \text{in } Q := (0, T) \times \Omega, \\ u(0, \cdot) = 0, & \text{in } \Omega, \\ u = f, & \text{on } \Sigma := (0, T) \times \Gamma, \end{cases} \quad (1.1)$$

where Δ_A is the Laplace operator associated with the magnetic potential $A \in W^{1,\infty}(\Omega)^3$,

$$\Delta_A := \sum_{j=1}^3 (\partial_{x_j} + ia_j)^2 = \Delta + 2iA \cdot \nabla + i(\nabla \cdot A) - |A|^2, \quad (1.2)$$

and $q \in L^\infty(\Omega)$. We define the Dirichlet-to-Neumann (DN) map associated with (1.1), as

$$\Lambda_{A,q}(f) := (\partial_\nu + iA \cdot \nu)u, \quad f \in L^2(\Sigma), \quad (1.3)$$

where $\nu(x)$ denotes the unit outward normal to $\partial\Omega$ at x , and u is the solution to (1.1).

In the remaining part of this text, two magnetic potentials $A_j \in W^{1,\infty}(\Omega)^3$, $j = 1, 2$, are said gauge equivalent, if there exists $\Psi \in W^{2,\infty}(\Omega)$ obeying $\Psi|_\Gamma = 0$, such that

$$A_2 = A_1 + \nabla \Psi. \quad (1.4)$$

In this paper we examine the uniqueness and stability issues in the inverse problem of determining the electric potential q and the gauge class of A , from the knowledge of $\Lambda_{A,q}$.

1.2. Physical motivations. The system (1.1) describes the quantum motion of a charged particle (the various physical constants are taken equal to 1) constrained by the unbounded domain Ω , under the influence of the magnetic field generated by A , and the electric potential q . Carbon nanotubes whose length-to-diameter ratio is up to $10^8/1$, are commonly modeled by infinite waveguides such as Ω . In this context, the inverse problem under consideration in this paper can be rephrased as to whether the strength of the electromagnetic

quantum disorder (namely, the magnetic field and the electric impurity potential q , see e.g. [21, 29]) can be determined by boundary measurement of the wave function u .

1.3. State of the art. Inverse coefficients problems for partial differential equations such as the Schrödinger equation, are the source of challenging mathematical problems, and have attracted many attention over the last decades. For instance, using the Bukhgeim-Klibanov method (see [14, 35, 36]), [3] claims Lipschitz stable determination of the time-independent electric potential perturbing the dynamic (i.e. non stationary) Schrödinger equation, from a single boundary measurement of the solution. In this case, the observation is performed on a sub boundary fulfilling the geometric optics condition for the observability, derived by Bardos, Lebeau and Rauch in [2]. This geometrical condition was removed by [8] for potentials which are *a priori* known in a neighborhood of the boundary, at the expense of weaker stability. In the same spirit, [22] Lipschitz stably determines by means of the Bukhgeim-Klibanov technique, the magnetic potential in the Coulomb gauge class, from a finite number of boundary measurements of the solution. Uniqueness results in inverse problems for the DN map related to the magnetic Schrödinger equation are also available in [24], but they are based on a different approach involving geometric optics (GO) solutions. The stable recovery of the magnetic field by the DN map of the dynamic magnetic Schrödinger equation is established in [9] by combining the approach used for determining the potential in hyperbolic equations (see [5, 7, 11, 28, 44, 47, 49]) with the one employed for the identification of the magnetic field in elliptic equations (see [23, 45, 50]). Notice that in the one-dimensional case, [1] proved by means of the boundary control method introduced by [4], that the DN map uniquely determines the time-independent electric potential of the Schrödinger equation. In [10] the time-independent electric potential is stably determined by the DN map associated with the dynamic magnetic Schrödinger equation on a Riemannian manifold. This result was recently extended by [6] to simultaneous determination of both the magnetic field and the electric potential. As for inverse coefficients problems of the Schrödinger equation with either Neumann, spectral, or scattering data, we refer to [23, 25, 26, 32, 37, 38, 45, 46, 50, 53].

All the above mentioned results are obtained in a bounded domain. Actually, there is only a small number of mathematical papers dealing with inverse coefficients problems in unbounded domains. One of them, [43], examines the problem of determining a potential appearing in the wave equation in the half-space. Assuming that the potential is known outside a fixed compact set, the author proves that it is uniquely determined by the DN map. Unique determination of compactly supported potentials appearing in the stationary Schrödinger equation in an infinite slab from partial DN measurements is established in [39]. The same problem is addressed by [37] for the stationary magnetic Schrödinger equation, and by [54] for bi-harmonic operators with perturbations of order zero or one. The inverse problem of determining the twisting function of an infinite twisted waveguide by the DN map, is addressed in [17]. The analysis carried out in [28, 44, 47, 49] is adapted to unbounded cylindrical domains in [17] for the determination of time-independent potentials with prescribed behavior outside a compact set, by the hyperbolic DN map. In [34], electric potentials with suitable exponential decay along the infinite direction of the waveguide, are stably recovered from a single boundary measurement of the solution. This is by means of a specifically designed Carleman estimate for the dynamic Schrödinger equation in infinite cylindrical domains, derived in [33]. The geometrical condition satisfied by the boundary data measurements in [34] is relaxed in [12] for potentials which are known in a neighborhood of the boundary. In [18], time-dependent potentials that are periodic in the translational direction of the waveguide, are stably retrieved by the DN map of the Schrödinger equation. In [30], periodic potentials are stably retrieved from the asymptotics of the boundary spectral data of the Dirichlet Laplacian. Finally, we refer to [19, 20], for the analysis of the Calderón problem in a periodic waveguide.

1.4. Well-posedness. We start by examining the well-posedness of the IBVP (1.1) in the functional space $\mathcal{C}([0, T], H^1(\Omega)) \cap \mathcal{C}^1([0, T], H^{-1}(\Omega))$. Namely, we are aiming for sufficient conditions on the coefficients

A , q and the non-homogeneous Dirichlet data f , ensuring that (1.1) admits a unique solution in the transposition sense. We say that $u \in L^\infty(0, T; H^{-1}(\Omega))$ is a solution to (1.1) in the transposition sense, if the identity

$$\langle u, F \rangle_{L^\infty(0, T; H^{-1}(\Omega)), L^1(0, T; H_0^1(\Omega))} = \langle f, \partial_\nu v \rangle_{L^2(\Sigma)},$$

holds for any $F \in L^1(0, T; H_0^1(\Omega))$. Here v denotes the unique $\mathcal{C}([0, T], H^1(\Omega))$ -solution to the transposition system

$$\begin{cases} (i\partial_t v + \Delta_A + q)v = F, & \text{in } Q, \\ v(T, \cdot) = 0, & \text{in } \Omega, \\ v = 0, & \text{on } \Sigma. \end{cases} \quad (1.5)$$

We refer to Subsection 2.3 for the full definition and description of transposition solutions to (1.1).

Since $\partial\Omega$ is not bounded, we introduce the following notations. First, we set

$$H^s(\partial\Omega) := H_{x_3}^s(\mathbb{R}, L^2(\partial\omega)) \cap L_{x_3}^2(\mathbb{R}, H^s(\partial\omega)), \quad s > 0,$$

where x_3 denotes the longitudinal variable of Ω . Next, we put

$$H^{r,s}((0, T) \times X) := H^r(0, T; L^2(X)) \cap L^2(0, T; H^s(X)), \quad r, s > 0,$$

where X is either Ω or $\partial\Omega$. For the sake of shortness, we write $H^{r,s}(Q)$ (resp., $H^{r,s}(\Sigma)$) instead of $H^{r,s}((0, T) \times \Omega)$ (resp., $H^{r,s}((0, T) \times \partial\Omega)$). Finally, we define

$$H_0^{2,1}(\Sigma) := \{f \in H^{2,1}(\Sigma); f(0, \cdot) = \partial_t f(0, \cdot) = 0\},$$

and state the existence and uniqueness result of solutions to (1.1) in the transposition sense, as follows.

Theorem 1.1. *For $M > 0$, let $A \in W^{1,\infty}(\Omega, \mathbb{R})^3$ and $q \in W^{1,\infty}(\Omega, \mathbb{R})$ satisfy the condition*

$$\|A\|_{W^{1,\infty}(\Omega)^3} + \|q\|_{W^{1,\infty}(\Omega)} \leq M. \quad (1.6)$$

Then, for each $f \in H_0^{2,1}(\Sigma)$, the IBVP (1.1) admits a unique solution in the transposition sense $u \in H^1(0, T; H^1(\Omega))$, and the estimate

$$\|u\|_{H^1(0, T; H^1(\Omega))} \leq C \|f\|_{H^{2,1}(\Sigma)}, \quad (1.7)$$

holds for some positive constant C depending only on T , ω and M . Moreover, the normal derivative $\partial_\nu u \in L^2(\Sigma)$, and we have

$$\|\partial_\nu u\|_{L^2(\Sigma)} \leq C \|f\|_{H^{2,1}(\Sigma)}. \quad (1.8)$$

It is clear from the definition (1.3) and the continuity property (1.8), that the DN map $\Lambda_{A,q}$ belongs to $\mathcal{B}(H_0^{2,1}(\Sigma), L^2(\Sigma))$, the set of linear bounded operators from $H_0^{2,1}(\Sigma)$ into $L^2(\Sigma)$.

1.5. Non uniqueness. There is a natural obstruction to the identification of A by $\Lambda_{A,q}$, arising from the invariance of the DN map under gauge transformation. More precisely, if $\Psi \in W^{2,\infty}(\Omega)$ verifies $\Psi|_\Gamma = 0$, then we have $u_{A+\nabla\Psi} = e^{-i\Psi} u_A$, where u_A (resp., $u_{A+\nabla\Psi}$) denotes the solution to (1.1) associated with the magnetic potential A (resp., $A + \nabla\Psi$), $q \in L^\infty(\Omega)$ and $f \in H_0^{2,1}(\Sigma)$. Further, as

$$(\partial_\nu + i(A + \nabla\Psi) \cdot \nu) u_{A+\nabla\Psi} = e^{-i\Psi} (\partial_\nu + iA \cdot \nu) u_A = (\partial_\nu + iA \cdot \nu) u_A \text{ on } \Sigma,$$

by direct calculation, we get that $\Lambda_{A,q} = \Lambda_{A+\nabla\Psi,q}$, despite of the fact that the two potentials A and $A + \nabla\Psi$ do not coincide in Ω (unless ψ is uniformly zero).

This shows that the best we can expect from the knowledge of the DN map is to identify (A, q) modulo gauge transformation of A . When $A|_{\partial\Omega}$ is known, this may be equivalently reformulated as to whether the magnetic field defined by the 2-form

$$dA := \sum_{i,j=1}^3 (\partial_{x_j} a_i - \partial_{x_i} a_j) dx_j \wedge dx_i,$$

and the electric potential q , can be retrieved by $\Lambda_{A,q}$. This is the inverse problem that we examine in the remaining part of this text.

1.6. Main results. We define the set of admissible magnetic potentials as

$$\mathcal{A} := \{A = (a_i)_{1 \leq i \leq 3}; a_1, a_2 \in L_{x_3}^\infty(\mathbb{R}, H_0^2(\omega)) \cap W^{2,\infty}(\Omega) \text{ and } a_3 \in C^3(\overline{\Omega}) \text{ satisfies (1.9) - (1.10)}\},$$

where

$$\sup_{x \in \Omega} \left(\sum_{\alpha \in \mathbb{N}^3, |\alpha| \leq 3} \langle x_3 \rangle^d |\partial_x^\alpha a_3(x)| \right) < \infty \text{ for some } d > 1, \quad (1.9)$$

and

$$\partial_x^\alpha a_3(x) = 0, \quad x \in \partial\Omega, \quad \alpha \in \mathbb{N}^3 \text{ such that } |\alpha| \leq 2. \quad (1.10)$$

Here $H_0^2(\omega)$ denotes the closure of $C_0^\infty(\omega)$ in the $H^2(\omega)$ -topology, and $\langle x_3 \rangle := (1 + x_3^2)^{1/2}$.

The first result of this paper claims stable determination of the magnetic field dA and unique identification of electric potential q , from the knowledge of the full data, i.e. the DN map defined by (1.3), where both the Dirichlet and Neumann measurements are performed on the whole boundary Σ .

Theorem 1.2. Fix $A_* := (a_{i,*})_{1 \leq i \leq 3} \in W^{2,\infty}(\Omega, \mathbb{R})^3$, and for $j = 1, 2$, let $q_j \in W^{1,\infty}(\Omega)$, and $A_j := (a_{i,j})_{1 \leq i \leq 3} \in A_* + \mathcal{A}$, satisfy the condition:

$$\sum_{i=1}^2 \partial_{x_i} (\partial_{x_3} (a_{i,1} - a_{i,2}) - \partial_{x_i} (a_{3,1} - a_{3,2})) = 0, \text{ in } \Omega. \quad (1.11)$$

Then, $\Lambda_{A_1, q_1} = \Lambda_{A_2, q_2}$ yields $(dA_1, q_1) = (dA_2, q_2)$.

Assume moreover that the estimate

$$\sum_{j=1}^2 (\|A_j\|_{W^{2,\infty}(\Omega)} + \|q_j\|_{W^{1,\infty}(\Omega)} + \|e_j\|_{W^{3,\infty}(\Omega)}) + \|A_*\|_{W^{2,\infty}(\Omega)} \leq M, \quad (1.12)$$

holds for some $M > 0$, with

$$e_j(x', x_3) := \int_{-\infty}^{x_3} (a_{3,j}(x', y_3) - a_{3,*}(x', y_3)) dy_3, \quad (x', x_3) \in \Omega.$$

Then there exist two constants $\mu_0 \in (0, 1)$ and $C > 0$, both of them depending only on T, ω and M , such that we have

$$\|dA_1 - dA_2\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{\mu_0}. \quad (1.13)$$

In (1.13) and in the remaining part of this text, $\|\cdot\|$ denotes the usual norm in $\mathcal{B}(H^{2,1}(\Sigma), L^2(\Sigma))$. Notice that in Theorem 1.2 we make use of the full DN map, as the magnetic field dA and the electric potential q are recovered by observing the solution to (1.1) on the entire lateral boundary Σ . In this case we may consider general unknown coefficients, in the sense that the behavior of A and q with respect to the infinite variable is not prescribed (we only assume that these coefficients and their derivatives are uniformly bounded in Ω). In order to achieve the same result by measuring on a bounded subset of Σ only, we need some extra information on the behavior of the unknown coefficients with respect to x_3 . Namely, we impose that the

strength of the magnetic field generated by $A = (a_i)_{1 \leq i \leq 3}$, reaches its maximum in the bounded subset $(-r, r) \times \omega$ of Ω , for some fixed $r > 0$, i.e.

$$\|\partial_{x_i} a_j - \partial_{x_j} a_i\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} = \|\partial_{x_i} a_j - \partial_{x_j} a_i\|_{L_{x_3}^\infty(-r, r; L^2(\omega))}, \quad i, j = 1, 2, 3. \quad (1.14)$$

Thus, with reference to (1.14), we set $\Gamma_r := \partial\omega \times (-r, r)$, introduce the space

$$H_0^{2,1}((0, T) \times \Gamma_r) := \{f \in H^{2,1}(\Sigma); f(0, \cdot) = \partial_t f(0, \cdot) = 0 \text{ and } \text{supp } f \subset [0, T] \times \partial\omega \times [-r, r]\},$$

and define the partial DN map $\Lambda_{A,q,r}$, by

$$\Lambda_{A,q,r}(f) := (\partial_\nu + iA \cdot \nu)u|_{(0,T) \times \Gamma_r}, \quad f \in H_0^{2,1}((0, T) \times \Gamma_r),$$

where u denotes the solution to (1.1). The following result states for each $r > 0$, that the magnetic field induced by potentials belonging (up to an additive $W^{2,\infty}(\Omega, \mathbb{R})^3$ -term) to

$$\mathcal{A}_r := \{A = (a_i)_{1 \leq i \leq 3} \in \mathcal{A} \text{ satisfying (1.14)}\},$$

can be retrieved from the knowledge of the partial DN map $\Lambda_{A,q,r'}$, provided we have $r' > r$.

Theorem 1.3. *For $j = 1, 2$, let $q_j \in W^{1,\infty}(\Omega, \mathbb{R})$, and let $A_j \in W^{2,\infty}(\Omega, \mathbb{R})^3$ satisfy $A_1 - A_2 \in \mathcal{A}_r$, for some $r > 0$. Suppose that there exists $r' > r$, such that $\Lambda_{A_1, q_1, r'} = \Lambda_{A_2, q_2, r'}$. Then, we have $dA_1 = dA_2$. Furthermore, if*

$$\|q_1 - q_2\|_{L_{x_3}^\infty(\mathbb{R}, H^{-1}(\omega))} = \|q_1 - q_2\|_{L_{x_3}^\infty(-r, r; H^{-1}(\omega))},$$

we have in addition $q_1 = q_2$.

Assume moreover that (1.11)-(1.12) hold. Then, the estimate

$$\|dA_1 - dA_2\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))^3} \leq C \|\Lambda_{A_1, q_1, r'} - \Lambda_{A_2, q_2, r'}\|^{\mu_1}, \quad (1.15)$$

holds with two constants $C > 0$, and $\mu_1 \in (0, 1)$, depending only on T, ω, M, r and r' .

We stress out that Theorem 1.3 applies not only to magnetic (resp., electric) potentials A_j (resp., q_j), $j = 1, 2$, which coincide outside $\omega \times (-r, r)$, but to a fairly more general class of magnetic potentials, containing, e.g., $2r$ -periodic potentials with respect to x_3 . More generally, if $g \in W^{2,\infty}(\mathbb{R}, \mathbb{R}_+)$ (resp. $g \in W^{1,\infty}(\mathbb{R}, \mathbb{R}_+)$) is an even and non-increasing function in \mathbb{R}_+ , then it is easy to see that potentials of the form $g \times A_j$ (resp., $g \times q_j$), where A_j (resp., q_j) are suitable $2r$ -periodic magnetic (resp., electric) potentials with respect to x_3 , fulfill the conditions of Theorem 1.3.

Notice that the absence of stability for the electric potential q , manifested in both Theorems 1.2 and 1.3, arises from the infinite extension of the spatial domain Ω in the x_3 direction. Indeed, the usual derivation of a stability equality for q , from estimates such as (1.13) or (1.15), requires that the differential operator d be invertible in Ω . Such a property is true in bounded domains (see e.g. [53]), but, to the best of our knowledge, it is not known whether it can be extended to unbounded waveguides. One way to overcome this technical difficulty is to impose certain gauge condition on the magnetic potentials, by prescribing their divergence. In this case, we establish in Theorem 1.4, below, that the electric and magnetic potentials can be simultaneously and stably determined by the DN map.

1.6.1. Simultaneous stable recovery of magnetic and electric potentials. We first introduce the set of divergence free transverse magnetic potentials,

$$\mathcal{A}_0 := \{A = (a_1, a_2, 0); a_1, a_2 \in L_{x_3}^\infty(\mathbb{R}, H_0^2(\omega)) \cap W^{2,\infty}(\Omega), \partial_{x_1} a_1 + \partial_{x_2} a_2 = 0 \text{ in } \Omega\},$$

in such a way that we have $\nabla \cdot A = \nabla \cdot A_*$ for any $A \in A_* + \mathcal{A}_0$. Here $A^* \in W^{2,\infty}(\Omega)^3$ is an arbitrary fixed magnetic potential. Since identifying $A \in A_* + \mathcal{A}_0$ from the knowledge of the DN map, amounts to determining the magnetic field dA , we have the following result.

Theorem 1.4. *Let $M > 0$, and let $A_* \in W^{2,\infty}(\Omega, \mathbb{R})^3$. For $j = 1, 2$, let $q_j \in W^{1,\infty}(\Omega, \mathbb{R})$, and let $A_j \in A_* + \mathcal{A}_0$ satisfy (1.12). Then, there exist two constant $\mu_2 \in (0, 1)$ and $C = C(T, \omega, M) > 0$, such that we have*

$$\|A_1 - A_2\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))^3} + \|q_1 - q_2\|_{L_{x_3}^\infty(\mathbb{R}, H^{-1}(\omega))} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{\mu_2}. \quad (1.16)$$

Assume moreover that the two following conditions

$$\|A_1 - A_2\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))^3} = \|A_1 - A_2\|_{L_{x_3}^\infty(-r, r; L^2(\omega))^3}, \quad (1.17)$$

and

$$\|q_1 - q_2\|_{L_{x_3}^\infty(\mathbb{R}, H^{-1}(\omega))} = \|q_1 - q_2\|_{L_{x_3}^\infty(-r, r; H^{-1}(\omega))}, \quad (1.18)$$

hold simultaneously for some $r > 0$. Then, for each $r' > r$, we have

$$\|A_1 - A_2\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} + \|q_1 - q_2\|_{L_{x_3}^\infty(\mathbb{R}, H^{-1}(\omega))} \leq C \|\Lambda_{A_1, q_1, r'} - \Lambda_{A_2, q_2, r'}\|^{\mu_2}, \quad (1.19)$$

where C is a positive constant depending only on T, ω, M, r and r' .

1.6.2. Comments. The key ingredient in the analysis of the inverse problem under examination is a suitable set of GO solutions to the magnetic Schrödinger equation appearing in (1.1). These functions are specifically designed for the waveguide geometry of Ω , in such a way that the unknown coefficients can be recovered by a separation of variables argument. More precisely, we seek GO solutions that are functions of $x = (x', x_3) \in \Omega$, but where the transverse variable $x' \in \omega$ and the translational variable $x_3 \in \mathbb{R}$ are separated. This approach was already used in [31], for determining zero order unknown coefficients of the wave equation. Since we consider first order unknown coefficients in this paper, the main issue here is to take into account both the cylindrical shape of Ω and the presence of the magnetic potential, in the design of the GO solutions.

When the domain Ω is bounded, we know from [9] that the magnetic field dA is uniquely determined by the DN map associated with (1.1). The main achievement of the present paper is to extend the above statement to unbounded cylindrical domains. Actually, we also improve the results of [9] in two directions. First, we prove simultaneous determination of the magnetic field dA and the electric potential q . Second, the regularity condition imposed on admissible magnetic potentials entering the Schrödinger equation of (1.1), is weakened from $W^{3,\infty}(\Omega)$ to $W^{2,\infty}(\Omega)$.

To our best knowledge, this is the first mathematical paper claiming identification by boundary measurements, of non-compactly supported magnetic field and electric potential. Moreover, in contrast to the other works [12, 18, 33] dealing with the stability issue of inverse problems for the Schrödinger equation in an infinite cylindrical domain, available in the mathematics literature, here we no longer require that the various unknown coefficients be periodic, or decay exponentially fast, in the translational direction of the waveguide.

Finally, since the conditions (1.14) and (1.17)-(1.18) are imposed in $\omega \times (-r, r)$ only, and since the solution to (1.1) lives in the infinitely extended cylinder $(0, T) \times \Omega$, we point out that the results of Theorems 1.3 and 1.4 cannot be derived from similar statements derived in a bounded domain.

1.7. Outlines. The paper is organized as follows. In Section 2 we examine the forward problem associated with (1.1), by rigorously defining the transposition solutions to (1.1), and proving Theorem 1.1. In Section 3, we build the GO solutions to the Schrödinger equation appearing in (1.1), which are the key ingredient in the analysis of the inverse problem carried out in the two last sections of this paper. In Section 4, we estimate the X-ray transform of first-order partial derivatives of the transverse magnetic potential, and the Fourier transform of the aligned magnetic field, in terms of the DN map. Finally, Section 5 contains the proofs of Theorems 1.2, 1.3, and 1.4.

2. ANALYSIS OF THE FORWARD PROBLEM

In this section we study the forward problem associated with (1.1), that is, we prove the statement of Theorem 1.1. Although this problem is very well documented when Ω is bounded (see e.g. [9]), to our best knowledge, it cannot be directly derived from any published mathematical work in the framework of the unbounded waveguide Ω under consideration in this paper.

The proof of Theorem 1.1, which is presented in Subsection 2.4, deals with transposition solutions to (1.1), that are rigorously defined in Subsection 2.3. As a preliminary, we start by examining in Subsection 2.1, the elliptic part of the dynamic magnetic Schrödinger operator appearing in (1.1), and we establish an existence and uniqueness result for the corresponding system in Subsection 2.2.

2.1. Elliptic magnetic Schrödinger operator. For $A \in W^{1,\infty}(\Omega, \mathbb{R})^3$, we set $\nabla_A := \nabla + iA$, where iA denotes the multiplier by iA , and notice for all $u \in H^1(\Omega)$, that

$$|\nabla_A u(x)|^2 \geq (1 - \epsilon)|\nabla u(x)|^2 + (1 - \epsilon^{-1})|Au(x)|^2, \quad \epsilon > 0, \quad x \in \Omega. \quad (2.1)$$

Next, for $q \in L^\infty(\Omega; \mathbb{R})$, we introduce the sesquilinear form

$$\mathbf{h}_{A,q}(u, v) := \int_{\Omega} \nabla_A u(x) \cdot \overline{\nabla_A v(x)} dx - \int_{\Omega} q(x) u(x) \overline{v(x)} dx, \quad u, v \in \mathcal{D}(\mathbf{h}_{A,q}) := H_0^1(\Omega),$$

and consider the self-adjoint operator $\mathcal{H}_{A,q}$ in $L^2(\Omega)$, generated by $\mathbf{h}_{A,q}$. In light of [34, Proposition 2.5], $\mathcal{H}_{A,q}$ acts on its domain $\mathcal{D}(\mathcal{H}_{A,q}) := H_0^1(\Omega) \cap H^2(\Omega)$, as the operator $-(\Delta_A + q)$, where $\Delta_A := \nabla_A \cdot \nabla_A$ is expressed by (1.2).

Further, for all $x \in \Omega$ fixed, taking $\epsilon = |A(x)|^2/(1 + |A(x)|^2)$ in (2.1), we get that $|\nabla_A u(x)|^2 \geq |\nabla u(x)|^2/(1 + |A(x)|^2) - |u(x)|^2$, whence

$$\mathbf{h}_{A,0}(u, u) + \|u\|_{L^2(\Omega)}^2 \geq \frac{\|\nabla u\|_{L^2(\Omega)}^2}{1 + \|A\|_{L^\infty(\Omega)}^2}, \quad u \in H_0^1(\Omega),$$

where $\mathbf{h}_{A,0}$ stands for $\mathbf{h}_{A,q}$ when q is uniformly zero. Thus, we deduce from the Poincaré inequality and Lax Milgram's theorem, that for any $v \in H^{-1}(\Omega)$, there exists a unique $\phi_v \in H_0^1(\Omega)$ satisfying

$$-\Delta_A \phi_v + \phi_v = v. \quad (2.2)$$

Next, for u and v in $H^{-1}(\Omega)$, we put

$$\langle u, v \rangle_{-1} := \operatorname{Re} \left(\int_{\Omega} \nabla_A \phi_u(x) \cdot \overline{\nabla_A \phi_v(x)} dx + \int_{\Omega} \phi_u(x) \overline{\phi_v(x)} dx \right),$$

and check that the space $H^{-1}(\Omega)$, endowed with the above scalar product, is Hilbertian. Having said that, we may now prove the following technical result.

Lemma 2.1. *For each $A \in W^{1,\infty}(\Omega, \mathbb{R})^3$, the linear operator $\mathcal{B}_A := \Delta_A$, with domain $\mathcal{D}(\mathcal{B}_A) := H_0^1(\Omega)$, is self-adjoint and negative in $H^{-1}(\Omega)$.*

Proof. We proceed as in the proof of [16, Proposition 2.6.14 and Corollary 2.6.15]. Namely, we pick u and v in $C_0^\infty(\Omega)$, and write

$$\langle \mathcal{B}_A u, v \rangle_{-1} = \langle w, v \rangle_{-1} + \langle u, v \rangle_{-1},$$

with $w := \mathcal{B}_A u - u$. Taking into account that $\phi_w = -u$, we obtain that

$$\langle \mathcal{B}_A u, v \rangle_{-1} = -\operatorname{Re} \left(\int_{\Omega} \nabla_A u(x) \cdot \overline{\nabla_A \phi_v(x)} dx + \int_{\Omega} u(x) \overline{\phi_v(x)} dx \right) + \langle u, v \rangle_{-1}. \quad (2.3)$$

Next, integrating by parts, we get that

$$-\operatorname{Re} \left(\int_{\Omega} \nabla_A u(x) \cdot \overline{\nabla_A \phi_v(x)} dx + \int_{\Omega} u(x) \overline{\phi_v(x)} dx \right) = -\operatorname{Re} \langle u, -\Delta_A \phi_v + \phi_v \rangle_{L^2(\Omega)} = -\operatorname{Re} \langle u, v \rangle_{L^2(\Omega)},$$

so (2.3) yields

$$\langle \mathcal{B}_A u, v \rangle_{-1} = -\operatorname{Re} \langle u, v \rangle_{L^2(\Omega)} + \langle u, v \rangle_{-1}. \quad (2.4)$$

Further, since $\langle u, u \rangle_{-1} = \operatorname{Re} \langle \phi_u, (-\Delta_A + 1) \phi_u \rangle_{L^2(\Omega)} = \operatorname{Re} \langle \phi_u, u \rangle_{L^2(\Omega)}$ and $\|\phi_u\|_{L^2(\Omega)}^2 \leq \langle u, u \rangle_{-1}$, we see that $\langle u, u \rangle_{-1} \leq \|u\|_{L^2(\Omega)}^2$. Therefore, we obtain

$$\langle \mathcal{B}_A u, u \rangle_{-1} = -\|u\|_{L^2(\Omega)}^2 + \langle u, u \rangle_{-1} \leq 0, \quad (2.5)$$

by taking $v = u$ in (2.4).

By density of $\mathcal{C}_0^\infty(\Omega)$ in $H_0^1(\Omega)$, both estimates (2.4) and (2.5) remain valid for all u and v in $H_0^1(\Omega)$. As a consequence, the operator \mathcal{B}_A is dissipative. Furthermore, $1 - \mathcal{B}_A$ being surjective from $H_0^1(\Omega)$ onto $H^{-1}(\Omega)$, by (2.2), we get that \mathcal{B}_A is m -dissipative. Moreover, as it follows readily from (2.4) that

$$\langle \mathcal{B}_A u, v \rangle_{-1} = \langle u, \mathcal{B}_A v \rangle_{-1}, \quad u, v \in H_0^1(\Omega),$$

we see that the graph of \mathcal{B}_A is contained into the one of its adjoint \mathcal{B}_A^* . Therefore, \mathcal{B}_A is self-adjoint, in virtue of [16, Corollary 2.4.10]. \square

2.2. Existence and uniqueness result. For further use, we establish the following existence and uniqueness result for the system

$$\begin{cases} (i\partial_t + \Delta_A + q)v = F, & \text{in } Q, \\ v(0, \cdot) = 0, & \text{in } \Omega, \\ v = 0, & \text{on } \Sigma, \end{cases} \quad (2.6)$$

with homogeneous Dirichlet boundary condition and suitable source term F .

Lemma 2.2. *Let M , A and q be the same as in Theorem 1.1.*

(i) *Assume that $F \in L^1(0, T; H_0^1(\Omega))$. Then, the system (2.6) admits a unique solution $v \in \mathcal{C}([0, T], H_0^1(\Omega))$, satisfying*

$$\|v\|_{\mathcal{C}([0, T], H^1(\Omega))} \leq C \|F\|_{L^1(0, T; H^1(\Omega))}, \quad (2.7)$$

for some constant $C > 0$, depending only on T , ω and M .

(ii) *If $F \in W^{1,1}(0, T; L^2(\Omega))$, then (2.6) admits a unique solution*

$$v \in \mathcal{Z} := \mathcal{C}^1([0, T], L^2(\Omega)) \cap \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega)),$$

and there exists $C = C(T, \omega, M) > 0$, such that

$$\|v\|_{\mathcal{Z}} \leq C \|F\|_{W^{1,1}(0, T; L^2(\Omega))}. \quad (2.8)$$

Proof. The proof boils down on the following statement, borrowed from [18, Lemma 2.1].

Let X be a Banach space, U be a m -dissipative operator in X with dense domain $D(U)$ and $B \in \mathcal{C}([0, T], \mathcal{B}(D(U)))$. Then for all $v_0 \in D(U)$ and $f \in \mathcal{C}([0, T], X) \cap L^1(0, T; D(U))$ (resp. $f \in W^{1,1}(0, T; X)$) there is a unique solution $v \in Z_0 = \mathcal{C}([0, T], D(U)) \cap C^1([0, T], X)$ to the following Cauchy problem

$$\begin{cases} v'(t) = Uv(t) + B(t)v(t) + f(t), \\ v(0) = v_0, \end{cases}$$

such that

$$\|v\|_{Z_0} = \|v\|_{C^0([0,T],D(U))} + \|v\|_{C^1([0,T],X)} \leq C(\|v_0\|_{D(U)} + \|f\|_*).$$

Here C is some positive constant depending only on T and $\|B\|_{C([0,T],\mathcal{B}(D(U)))}$, and $\|f\|_*$ stands for the norm $\|f\|_{C([0,T],X) \cap L^1(0,T;D(U))}$ (resp. $\|f\|_{W^{1,1}(0,T;X)}$).

Notice that the operator $i\mathcal{B}_A$ is skew-adjoint, since \mathcal{B}_A is self-adjoint in $H^{-1}(\Omega)$. Hence $i\mathcal{B}_A$ is m -dissipative with dense domain in $H^{-1}(\Omega)$. Further, the multiplier by iq being bounded in $\mathcal{C}[0,T]$, $H_0^1(\Omega)$, we obtain (i) by applying the above result with $X = H^{-1}(\Omega)$, $U = i\mathcal{B}_{A,q}$, $f = iF$, $B(t) = iq$, and $v_0 = 0$.

Similarly, as $\mathcal{H}_{A,q}$ is self-adjoint in $L^2(\Omega)$, then the operator $-i\mathcal{H}_{A,q}$ is m -dissipative with dense domain in $L^2(\Omega)$, so we derive (ii) by applying [18, Lemma 2.1] with $X = L^2(\Omega)$, $U = -i\mathcal{H}_{A,q}$, $f = iF$, $B(t) = 0$, and $v_0 = 0$. \square

Remark 2.3. Since $w(t, x) := \overline{v(T - t, x)}$, for $(t, x) \in Q$, is solution to

$$\begin{cases} (i\partial_t + \Delta_A + q)w = F, & \text{in } Q, \\ w(T, \cdot) = 0, & \text{in } \Omega, \\ w = 0 & \text{on } \Sigma, \end{cases} \quad (2.9)$$

whenever v is solution to the IBVP (2.6), where the function $(t, x) \mapsto \overline{v(T - t, x)}$ is substituted for F , we infer from Lemma 2.2 that the transposed system (2.9) admits a unique solution w in $C^0([0, T], H_0^1(\Omega))$ (resp., \mathcal{Z}) provided F is in $L^1(0, T; H_0^1(\Omega))$ (resp., $W^{1,1}(0, T; L^2(\Omega))$).

2.3. Transposition solutions. As a preamble to the definition of transposition solutions to (1.1), we establish that the normal derivative of the $\mathcal{C}([0, T], H_0^1(\Omega))$ -solution to (2.6) is lying in $L^2(\Sigma)$.

Lemma 2.4. Let M , A and q be as in Lemma 2.2. Then, the linear map $F \mapsto \partial_\nu v$, where v denotes the $\mathcal{C}([0, T], H_0^1(\Omega))$ -solution to (2.6) associated with $F \in L^1(0, T; H_0^1(\Omega))$, given by Lemma 2.2, is bounded from $L^1(0, T; H_0^1(\Omega))$ into $L^2(\Sigma)$.

Proof. Since $\|v\|_{\mathcal{C}([0,T],H^1(\Omega))} \leq C\|F\|_{L^1(0,T;H^1(\Omega))}$, by (2.7), we may assume without loss of generality that $A = 0$ and $q = 0$.

Assume that $F \in W^{1,1}(0, T; L^2(\Omega))$ in such a way $v \in \mathcal{Z}$, in virtue of Lemma 2.2. Let $N_1 \in \mathcal{C}^2(\overline{\omega})^2$ satisfy $N_1 = \nu_1$ on $\partial\omega$, where ν_1 denotes the unit outward normal vector to $\partial\omega$. Put $N(x', x_3) := (N_1(x'), 0)$ for all $x' \in \omega$ and $x_3 \in \mathbb{R}$, so that $N \in \mathcal{C}^2(\overline{\Omega})^3 \cap W^{2,\infty}(\Omega)^3$ verifies $N = \nu$ on $\partial\Omega$. Then, we have

$$\langle i\partial_t v + \Delta v, N \cdot \nabla v \rangle_{L^2(Q)} = \langle F, N \cdot \nabla v \rangle_{L^2(Q)}. \quad (2.10)$$

By integrating by parts with respect to t , we get

$$\begin{aligned} \langle \partial_t v, N \cdot \nabla v \rangle_{L^2(Q)} &= \langle v(T, \cdot), N \cdot \nabla v(T, \cdot) \rangle_{L^2(\Omega)} - \langle v, N \cdot \nabla \partial_t v \rangle_{L^2(Q)} \\ &= \langle v(T, \cdot), N \cdot \nabla v(T, \cdot) \rangle_{L^2(\Omega)} + \langle N \cdot \nabla v, \partial_t v \rangle_{L^2(Q)} - I, \end{aligned} \quad (2.11)$$

where $I := \int_Q N \cdot \nabla(v \overline{\partial_t v}) dx dt$. Taking into account that $N \cdot \nabla = N_1 \cdot \nabla_{x'}$, where $\nabla_{x'}$ denotes the gradient operator with respect to $x' \in \omega$, we have $I = \int_Q N_1 \cdot \nabla_{x'}(v \overline{\partial_t v}) dx dt$, hence

$$\begin{aligned} I &= \int_Q \nabla_{x'} \cdot (v(t, x) \overline{\partial_t v}(t, x) N_1(x')) dx' dx_3 dt - \langle (\nabla \cdot N) v, \overline{\partial_t v} \rangle_{L^2(Q)} \\ &= \int_\Sigma v(t, x) \overline{\partial_t v}(t, x) N_1(x') \cdot \nu_1(x') dx' dx_3 dt - \langle (\nabla \cdot N) v, \partial_t v \rangle_{L^2(Q)} \\ &= -\langle (\nabla \cdot N) v, \partial_t v \rangle_{L^2(Q)}, \end{aligned} \quad (2.12)$$

by Green's formula, since $v|_{\Sigma} = 0$. Putting (2.11)-(2.12) together, we obtain that

$$\begin{aligned} & 2\operatorname{Re}\langle i\partial_t v, N \cdot \nabla v \rangle_{L^2(Q)} \\ &= i\langle v(T, \cdot), N \cdot \nabla v(T, \cdot) \rangle_{L^2(\Omega)} - \langle (\nabla \cdot N)v, i\partial_t v \rangle_{L^2(Q)} \\ &= i\langle v(T, \cdot), N \cdot \nabla v(T, \cdot) \rangle_{L^2(\Omega)} + \langle (\nabla \cdot N)v, \Delta v \rangle_{L^2(Q)} - \langle (\nabla \cdot N)v, F \rangle_{L^2(Q)}. \end{aligned} \quad (2.13)$$

Applying the Green formula with respect to $x' \in \omega$ and integrating by parts with respect to $x_3 \in \mathbb{R}$, we find that

$$\langle (\nabla \cdot N)v, \Delta v \rangle_{L^2(Q)} = -\langle (\nabla \cdot N)\nabla v, \nabla v \rangle_{L^2(Q)^3} - \langle v\nabla(\nabla \cdot N), \nabla v \rangle_{L^2(Q)^3},$$

so (2.13) entails

$$\begin{aligned} 2\operatorname{Re}\langle i\partial_t v, N \cdot \nabla v \rangle_{L^2(Q)} &= i\langle v(T, \cdot), N \cdot \nabla v(T, \cdot) \rangle_{L^2(\Omega)} - \langle (\nabla \cdot N)\nabla v, \nabla v \rangle_{L^2(Q)^3} \\ &\quad - \langle v\nabla(\nabla \cdot N), \nabla v \rangle_{L^2(Q)^3} - \langle (\nabla \cdot N)v, F \rangle_{L^2(Q)}. \end{aligned}$$

This and (2.7) yield

$$\begin{aligned} |\operatorname{Re}\langle i\partial_t v, N \cdot \nabla v \rangle_{L^2(Q)}| &\leq C\|v\|_{C([0,T],H^1(\Omega))} (\|v\|_{C([0,T],H^1(\Omega))} + \|F\|_{L^1(0,T;H^1(\Omega))}) \\ &\leq C\|F\|_{L^1(0,T;H^1(\Omega))}^2. \end{aligned}$$

From this and (2.10), it then follows that

$$|\operatorname{Re}\langle \Delta v, N \cdot \nabla v \rangle_{L^2(Q)}| \leq C\|F\|_{L^1(0,T;H^1(\Omega))}^2 \quad (2.14)$$

On the other hand, we get upon applying the Green formula with respect to $x' \in \omega$ and integrating by parts with respect to $x_3 \in \mathbb{R}$, that

$$\begin{aligned} \langle \Delta v, N \cdot \nabla v \rangle_{L^2(Q)} &= -\langle \nabla v, \nabla(N \cdot \nabla v) \rangle_{L^2(Q)^3} + \langle \nabla v \cdot \nu, N \cdot \nabla v \rangle_{L^2(\Sigma)} \\ &= -\langle \nabla v, \nabla(N \cdot \nabla v) \rangle_{L^2(Q)^3} + \|\partial_\nu v\|_{L^2(\Sigma)}^2. \end{aligned} \quad (2.15)$$

Moreover, since $\operatorname{Re}(\nabla v \cdot \nabla(N \cdot \overline{\nabla v})) = \operatorname{Re}((H\nabla v) \cdot \overline{\nabla v}) + \frac{1}{2}N \cdot \nabla |\nabla v|^2$ with $H := (\partial_{x_i} N_j)_{1 \leq i,j \leq 3}$ and $N := (N_j)_{1 \leq j \leq 3}$, we infer from (2.15) that

$$\operatorname{Re}\langle \Delta v, N \cdot \nabla v \rangle_{L^2(Q)} = \|\partial_\nu v\|_{L^2(\Sigma)}^2 - \operatorname{Re}\langle H\nabla v, \nabla v \rangle_{L^2(Q)^3} - \frac{1}{2} \int_Q N \cdot \nabla |\nabla v|^2 \, dx dt. \quad (2.16)$$

Further, by applying once more the Green formula with respect to $x' \in \omega$, we find for a.e. $(t, x_3) \in (0, T) \times \mathbb{R}$, that

$$\begin{aligned} & \int_\omega N(x', x_3) \cdot \nabla |\nabla v(t, x', x_3)|^2 \, dx' = \int_\omega N_1(x') \cdot \nabla_{x'} |\nabla v(t, x', x_3)|^2 \, dx' \\ &= \|\nabla v(t, \cdot, x_3)\|_{L^2(\partial\omega)^3}^2 - \langle (\nabla \cdot N)\nabla v(t, \cdot, x_3), \nabla v(t, \cdot, x_3) \rangle_{L^2(\omega)^3}. \end{aligned} \quad (2.17)$$

Bearing in mind that $v|_{\Sigma} = 0$, we have $|\nabla v|^2 = |\partial_\nu v|^2$ on Σ , so we deduce from (2.17) that

$$\int_Q N \cdot \nabla |\nabla v|^2 \, dx dt = \|\partial_\nu v\|_{L^2(\Sigma)}^2 - \langle (\nabla \cdot N)\nabla v, \nabla v \rangle_{L^2(Q)^3}.$$

From this and (2.16), it then follows that

$$\|\partial_\nu v\|_{L^2(\Sigma)}^2 = 2\operatorname{Re}\langle \Delta v, N \cdot \nabla v \rangle_{L^2(Q)} + 2\operatorname{Re}\langle H\nabla v, \nabla v \rangle_{L^2(Q)^3} - \langle (\nabla \cdot N)\nabla v, \nabla v \rangle_{L^2(Q)^3},$$

and hence

$$\|\partial_\nu v\|_{L^2(\Sigma)} \leq C(\|F\|_{L^1(0,T;H^1(\Omega))} + \|v\|_{C([0,T],H^1(\Omega))}) \leq C\|F\|_{L^1(0,T;H^1(\Omega))},$$

according to (2.7) and (2.14). By density of $W^{1,1}(0, T; H_0^1(\Omega))$ in $L^1(0, T; H_0^1(\Omega))$, it is clear that the above estimate extends to every $F \in L^1(0, T; H_0^1(\Omega))$, which proves the desired result. \square

Armed with Lemma 2.4, we may now introduce the transposition solution to (1.1). For $F \in L^1(0, T; H_0^1(\Omega))$, we denote by $v \in C^0([0, T], H_0^1(\Omega))$ the solution to (2.9), given by Remark 2.3. Since $(t, x) \mapsto \overline{v(T-t, x)}$ is solution to (2.6) associated with the source term $(t, x) \mapsto \overline{F(T-t, x)}$, we infer from Lemma 2.4, that the mapping $F \mapsto \partial_\nu v$ is bounded from $L^1(0, T; H_0^1(\Omega))$ into $L^2(\Sigma)$. Therefore, for each $f \in L^2(\Sigma)$, the mapping

$$\ell_f : F \mapsto \langle f, \partial_\nu v \rangle_{L^2(\Sigma)},$$

is an anti-linear form on $L^1(0, T; H_0^1(\Omega))$. Thus, there exists a unique $u \in L^\infty(0, T; H^{-1}(\Omega))$ such that we have

$$\langle u, F \rangle_{L^\infty(0, T; H^{-1}(\Omega)), L^1(0, T; H_0^1(\Omega))} = \ell_f(F), \quad F \in L^1(0, T; H_0^1(\Omega)), \quad (2.18)$$

according to Riesz's representation theorem. The function u , characterized by (2.18), is named the solution in the transposition sense to (1.1).

2.4. Proof of Theorem 1.1. Let $w \in L^\infty(0, T; H^{-1}(\Omega))$ be the solution in the transposition sense to the system

$$\begin{cases} (i\partial_t + \Delta_A + q)w = 0, & \text{in } Q, \\ w(0, \cdot) = 0, & \text{in } \Omega, \\ w = \partial_t^2 f, & \text{on } \Sigma. \end{cases}$$

For any $t \in (0, T)$, put $v(t, \cdot) := \int_0^t w(s, \cdot) ds$, in such a way that v is the solution in the transposition sense to the system

$$\begin{cases} (i\partial_t + \Delta_A + q)v = 0, & \text{in } Q, \\ v(0, \cdot) = 0, & \text{in } \Omega, \\ v = \partial_t f, & \text{on } \Sigma. \end{cases} \quad (2.19)$$

We have $v = \partial_t f \in H^{1,1/2}(\Sigma)$ by [41, Section 4, Proposition 2.3], and since $H^{1,1/2}(\Sigma) \subset L^2(0, T; H^{1/2}(\partial\Omega))$, and $-\Delta_A v = iw + qv$ in Q , from the first line of (2.19), then $v \in L^2(0, T; H^1(\Omega)) \cap W^{1,\infty}(0, T; H^{-1}(\Omega))$. Moreover, we have the following estimate

$$\|v\|_{L^2(0, T; H^1(\Omega))} \leq C \left(\|w\|_{L^2(0, T; H^{-1}(\Omega))} + \|qv\|_{L^2(0, T; H^{-1}(\Omega))} + \|\partial_t f\|_{L^2(0, T; H^{1/2}(\partial\Omega))} \right), \quad (2.20)$$

where the constant $C > 0$ depends only on T, ω , and M .

On the other hand, from the very definition of the transposition solution w , we obtain

$$\|w\|_{L^2(0, T; H^{-1}(\Omega))} \leq T^{1/2} \|w\|_{L^\infty(0, T; H^{-1}(\Omega))} \leq C \|\partial_t^2 f\|_{L^2(\Sigma)} \leq C \|f\|_{H^{2,1}(\Sigma)}, \quad (2.21)$$

with the aid of Lemma 2.4. As a consequence we have

$$\|qv\|_{L^2(0, T; H^{-1}(\Omega))} \leq \|q\|_{W^{1,\infty}(\Omega)} T \|w\|_{L^2(0, T; H^{-1}(\Omega))} \leq C \|f\|_{H^{2,1}(\Sigma)}. \quad (2.22)$$

Putting (2.20)–(2.22) together, we find that

$$\|v\|_{L^2(0, T; H^1(\Omega))} \leq C \|f\|_{H^{2,1}(\Sigma)}, \quad (2.23)$$

for some constant $C = C(T, \omega, M) > 0$.

Finally, as $u(t) = \int_0^t v(s) ds$ is solution to (1.1) in the transposition sense, we have

$$\|u\|_{H^1(0, T; H^1(\Omega))} \leq (1 + T)^{1/2} \|v\|_{L^2(0, T; H^1(\Omega))},$$

hence (1.7) follows from this and (2.23).

We turn now to proving (1.8). To do that, we pick $f \in \mathcal{C}^\infty([0, T] \times \partial\Omega) \cap H_0^{2,1}(\Sigma)$, and proceed as in the derivation of Lemma 2.4. We get that

$$\|\partial_\nu u\|_{L^2(\Sigma)} \leq C (\|u\|_{H^1(0,T;H^1(\Omega))} + \|f\|_{H^{2,1}(\Sigma)}),$$

for some constant $C = C(T, \omega, M) > 0$, so we deduce from (1.7) that

$$\|\partial_\nu u\|_{L^2(\Sigma)} \leq C \|f\|_{H^{2,1}(\Sigma)}.$$

The desired result follows from this by density of $\mathcal{C}^\infty([0, T] \times \partial\Omega) \cap H_0^{2,1}(\Sigma)$ in $H_0^{2,1}(\Sigma)$.

3. GO SOLUTIONS

In this section we build GO solutions to the magnetic Schrödinger equation in Ω . These functions are essential tools in the proof of Theorems 1.2, 1.3 and 1.4. As in [32], we take advantage of the translational invariance of Ω with respect to the longitudinal direction x_3 , in order to adapt the method suggested by Bellassoued and Choulli in [9] for building GO solutions to the magnetic Schrödinger equation in a bounded domain, to the framework of the unbounded waveguide Ω . Moreover, as we aim to reduce the regularity assumption imposed on the magnetic potential by the GO solutions construction method, we follow the strategy developed in [23, 37, 38, 45] for magnetic Laplace operators, and rather build GO solutions to the Schrödinger equation associated with a suitable smooth approximation of the magnetic potential.

Throughout the entire section, we consider two magnetic potentials

$$A_j = (A_j^\sharp, a_{j,3}) \in W^{2,\infty}(\Omega, \mathbb{R})^2 \times W^{2,\infty}(\Omega, \mathbb{R}), \quad j = 1, 2,$$

and two electric potentials $q_j \in W^{1,\infty}(\Omega, \mathbb{R})$, obeying the conditions

$$\|A_j\|_{W^{2,\infty}(\Omega)^3} + \|q_j\|_{W^{1,\infty}(\Omega)} \leq M, \quad j = 1, 2, \quad (3.1)$$

and

$$\partial_x^\alpha A_1 = \partial_x^\alpha A_2 \text{ on } \partial\Omega, \text{ for all } \alpha \in \mathbb{N}^3 \text{ such that } |\alpha| \leq 1. \quad (3.2)$$

For $\sigma > 0$, we denote by $A_{j,\sigma}^\sharp$ a suitable $\mathcal{C}^\infty(\mathbb{R}^3, \mathbb{R})^2 \cap W^{\infty,\infty}(\mathbb{R}^3, \mathbb{R})^2$ -approximation of A_j^\sharp , we shall specify in Lemma 3.3, below. We seek solutions $u_{j,\sigma}$ to the magnetic Schrödinger equation of (1.1), where (A_j, q_j) is substituted for (A, q) , of the form

$$u_{j,\sigma}(t, x', x_3) := \Phi_j(2\sigma t, x) b_{j,\sigma}(2\sigma t, x) e^{i\sigma(x' \cdot \theta - \sigma t)} + \psi_{j,\sigma}(t, x), \quad t \in \mathbb{R}, \quad x = (x', x_3) \in \omega \times \mathbb{R}. \quad (3.3)$$

Here, $\theta \in \mathbb{S}^1 := \{y \in \mathbb{R}^2 : |y| = 1\}$ is fixed,

$$b_{j,\sigma}(t, x) := \exp\left(-i \int_0^t \theta \cdot A_{j,\sigma}^\sharp(x' - s\theta, x_3) ds\right), \quad t \in \mathbb{R}, \quad x = (x', x_3) \in \omega \times \mathbb{R}, \quad (3.4)$$

Φ_j is a solution to the following transport equation

$$(\partial_t + \theta \cdot \nabla_{x'}) \Phi_j = 0 \text{ in } \mathbb{R} \times \Omega, \quad (3.5)$$

and we imposed that the remainder term $\psi_{j,\sigma} \in L^2(Q)$ scales at best like $\sigma^{-1/2}$ when σ is large, i.e.

$$\lim_{\sigma \rightarrow +\infty} \sigma^{1/2} \|\psi_{j,\sigma}\|_{L^2(Q)} = 0. \quad (3.6)$$

Such a construction requires that $A_{j,\sigma}^\sharp$ be sufficiently close to A_j^\sharp , as will appear in the coming subsection.

3.1. Magnetic potential mollification. We aim to define a suitable smooth approximation $A_{j,\sigma}^\# \in C^\infty(\mathbb{R}^3, \mathbb{R})^2 \cap W^{\infty,\infty}(\mathbb{R}^3, \mathbb{R})^2$ of $A_j^\# = (a_{1,j}, a_{2,j})$, for $j = 1, 2$. This preliminarily requires that $A_j^\#$ be appropriately extended to a larger domain than Ω , as follows.

Lemma 3.1. *Let $A_j^\#$, for $j = 1, 2$, be in $W^{2,\infty}(\Omega, \mathbb{R})^2$ and satisfy (3.2). Let $\tilde{\omega}$ be a smooth open bounded subset of \mathbb{R}^2 , containing $\bar{\omega}$. Then, there exist two potentials $\tilde{A}_1^\#$ and $\tilde{A}_2^\#$ in $W^{2,\infty}(\mathbb{R}^3, \mathbb{R})^2$, both of them being supported in $\tilde{\Omega} := \tilde{\omega} \times \mathbb{R}$, such that we have*

$$\tilde{A}_j^\# = A_j^\# \text{ in } \Omega, \text{ for } j = 1, 2, \text{ and } \tilde{A}_1^\# = \tilde{A}_2^\# \text{ in } \tilde{\Omega} \setminus \Omega. \quad (3.7)$$

Moreover, the two estimates

$$\|\tilde{A}_j^\#\|_{W^{2,\infty}(\mathbb{R}^3)^2} \leq C \max \left(\|A_1^\#\|_{W^{2,\infty}(\Omega)^2}, \|A_2^\#\|_{W^{2,\infty}(\Omega)^2} \right), \quad j = 1, 2, \quad (3.8)$$

hold for some constant $C > 0$, depending only on ω and $\tilde{\omega}$.

Proof. By [48, Section 3, Theorem 5] and [34, Lemma 2.7], there exists $\tilde{A}_1^\# \in W^{2,\infty}(\mathbb{R}^3, \mathbb{R})^2$, such that $\tilde{A}_1^\# = A_1^\#$ in Ω , and (3.8) holds true for $j = 1$. Then, upon possibly substituting $\chi \tilde{A}_1^\#$ for $\tilde{A}_1^\#$, where $\chi \in C_0^\infty(\mathbb{R}^3, \mathbb{R})$ is supported in $\tilde{\Omega}$ and verifies $\chi(x) = 1$ for all $x \in \Omega$, we may assume that $\tilde{A}_1^\#$ is supported in $\tilde{\Omega}$ as well.

Next, we put

$$\tilde{A}_2^\#(x) := \begin{cases} A_2^\#(x), & \text{if } x \in \Omega, \\ \tilde{A}_1^\#(x), & \text{if } x \in \mathbb{R}^3 \setminus \Omega. \end{cases} \quad (3.9)$$

Then, it is clear from (3.2) that $\tilde{A}_2^\# \in W^{2,\infty}(\mathbb{R}^3, \mathbb{R})^2$ and that it satisfies (3.8) with $j = 2$. \square

Having seen this, we define for each $\sigma > 0$ the smooth approximation $a_\sigma \in C^\infty(\mathbb{R}^3, \mathbb{R}) \cap W^{\infty,\infty}(\mathbb{R}^3, \mathbb{R})$ of a function $\tilde{a} \in W^{2,\infty}(\mathbb{R}^3, \mathbb{R})$, supported in $\tilde{\Omega}$, by

$$a_\sigma(x) := \int_{\mathbb{R}^3} \chi_\sigma(x-y) (\tilde{a}(y) + (x-y) \cdot \nabla \tilde{a}(y)) dy, \quad x \in \mathbb{R}^3. \quad (3.10)$$

Here we have set $\chi_\sigma(x) := \sigma \chi(\sigma^{1/3}x)$ for all $x \in \mathbb{R}^3$, where $\chi \in C_0^\infty(\mathbb{R}^3, \mathbb{R}_+)$ is such that

$$\text{supp } \chi \subset \{x \in \mathbb{R}^3; |x| \leq 1\} \text{ and } \int_{\mathbb{R}^3} \chi(x) dx = 1.$$

This terminology is justified by the fact that a_σ gets closer to \tilde{a} as the parameter σ becomes larger, as can be seen from the following result.

Lemma 3.2. *Let $\tilde{a} \in W^{2,\infty}(\mathbb{R}^3, \mathbb{R})$ be supported in $\tilde{\Omega}$ and satisfy $\|\tilde{a}\|_{W^{2,\infty}(\mathbb{R}^3)} \leq M$, for some $M > 0$. Then, there exists a constant $C > 0$, depending only on ω , $\tilde{\omega}$, and M , such that for all $\sigma > 0$, we have*

$$\|a_\sigma - \tilde{a}\|_{W^{k,\infty}(\mathbb{R}^3)} \leq C \sigma^{(k-2)/3}, \quad k = 0, 1, \quad (3.11)$$

where $W^{0,\infty}(\Omega)$ stands for $L^\infty(\Omega)$, and

$$\|a_\sigma\|_{W^{k,\infty}(\mathbb{R}^3)} \leq C \sigma^{(k-2)/3}, \quad k \geq 2. \quad (3.12)$$

Proof. We only establish (3.11), the estimate (3.12) being obtained with similar arguments. For $x \in \mathbb{R}^3$ fixed, we make the change of variable $\eta = \sigma^{1/3}(x-y)$ in (3.10). We get

$$a_\sigma(x) = \int_{\mathbb{R}^3} \chi(\eta) \tilde{a}(x - \sigma^{-1/3}\eta) d\eta + \sigma^{-1/3} \int_{\mathbb{R}^3} \chi(\eta) (\eta \cdot \nabla \tilde{a}(x - \sigma^{-1/3}\eta)) d\eta. \quad (3.13)$$

On the other hand, we have

$$\begin{aligned} \int_{\mathbb{R}^3} \chi(\eta) \tilde{a}(x - \sigma^{-1/3} \eta) d\eta - \tilde{a}(x) &= \int_{\mathbb{R}^3} \chi(\eta) \left(\tilde{a}(x - \sigma^{-1/3} \eta) - \tilde{a}(x) \right) d\eta \\ &= -\sigma^{-1/3} \int_{\mathbb{R}^3} \chi(\eta) \left(\int_0^1 \eta \cdot \nabla \tilde{a}(x - s\sigma^{-1/3} \eta) ds \right) d\eta, \end{aligned}$$

so we infer from (3.13) that

$$a_\sigma(x) - \tilde{a}(x) = \sigma^{-1/3} \int_{\mathbb{R}^3} \chi(\eta) \left(\int_0^1 \eta \cdot \left(\nabla \tilde{a}(x - \sigma^{-1/3} \eta) - \nabla \tilde{a}(x - s\sigma^{-1/3} \eta) \right) ds \right) d\eta. \quad (3.14)$$

By the Sobolev embedding theorem (see e.g. [27, Theorem 1.4.4.1] and [51, Lemma 3.13]), we know that $\tilde{a} \in C^{1,1}(\mathbb{R}^3)$ satisfies the estimate $\|\tilde{a}\|_{C^{1,1}(\mathbb{R}^3)} \leq C \|\tilde{a}\|_{W^{2,\infty}(\mathbb{R}^3)}$, where $C > 0$ is independent of \tilde{a} . From this and (3.14), it then follows that

$$|a_\sigma(x) - \tilde{a}(x)| \leq C \|\tilde{a}\|_{W^{2,\infty}(\mathbb{R}^3)} \left(\int_{\mathbb{R}^3} \chi(\eta) |\eta|^2 d\eta \right) \sigma^{-2/3},$$

which, together with the estimate $\|\tilde{a}\|_{W^{2,\infty}(\mathbb{R}^3)} \leq M$, yields (3.11) with $k = 0$. Further, upon differentiating (3.14) with respect to x_i , for $i = 1, 2, 3$, and upper bounding the integrand function $(\eta, s) \mapsto \nabla \partial_i \tilde{a}(x - \sigma^{-1/3} \eta) - \nabla \partial_i \tilde{a}(x - s\sigma^{-1/3} \eta)$ by $2\|\tilde{a}\|_{W^{2,\infty}(\mathbb{R}^3)}$, uniformly over $\mathbb{R}^3 \times (0, 1)$, we obtain (3.11) for $k = 1$. \square

We notice for further use from (3.10) and the expression of χ_σ , that

$$\begin{aligned} a_\sigma(x) &= \int_{\mathbb{R}^3} (\chi_\sigma(x - y) - \nabla \cdot ((x - y) \chi_\sigma(x - y))) \tilde{a}(y) dy \\ &= \int_{\mathbb{R}^3} \left(4\sigma \chi(\sigma^{1/3}(x - y)) + \sigma^{4/3}(x - y) \cdot \nabla \chi(\sigma^{1/3}(x - y)) \right) \tilde{a}(y) dy, \quad x \in \mathbb{R}^3. \end{aligned}$$

Making the change of variable $z = \sigma^{1/3}(x - y)$ in the above integral, we find that

$$a_\sigma(x) = \int_{\mathbb{R}^3} (4\chi(z) + z \cdot \nabla \chi(z)) \tilde{a}(\sigma^{-1/3}z - x) dz, \quad x \in \mathbb{R}^3.$$

Since χ is compactly supported in \mathbb{R}^3 , this entails that

$$\|a_\sigma\|_{L^\infty(\mathbb{R}^3)} \leq C \|\tilde{a}\|_{L^\infty(\mathbb{R}^3)}, \quad \sigma > 0, \quad (3.15)$$

where the constant $C > 0$ depends only on χ .

Let $\tilde{A}_j^\# = (\tilde{a}_{1,j}, \tilde{a}_{2,j})$, $j = 1, 2$, be given by Lemma 3.1. With reference to (3.10), we define the smooth magnetic potentials

$$A_{j,\sigma}^\# = (a_{1,j,\sigma}, a_{2,j,\sigma}) \in C^\infty(\mathbb{R}^3, \mathbb{R})^2 \cap W^{\infty,\infty}(\mathbb{R}^3, \mathbb{R})^2,$$

by setting

$$a_{i,j,\sigma}(x) := \int_{\mathbb{R}^3} \chi_\sigma(x - y) (\tilde{a}_{i,j}(y) + (x - y) \cdot \nabla \tilde{a}_{i,j}(y)) dy, \quad x \in \mathbb{R}^3, \quad i, j = 1, 2. \quad (3.16)$$

Thus, applying Lemma 3.2 with $\tilde{a} = \tilde{a}_{i,j}$ for $i, j = 1, 2$, we obtain the following result.

Lemma 3.3. *For $j = 1, 2$, let $A_j^\#$ be the same as in Lemma 3.1, and satisfy (3.1). Then, there exists a constant $C > 0$, depending only on ω and M , such that for each $j = 1, 2$, and all $\sigma > 0$, we have*

$$\|A_{j,\sigma}^\# - A_j^\#\|_{W^{k,\infty}(\Omega)^2} \leq \|A_{j,\sigma}^\# - \tilde{A}_j^\#\|_{W^{k,\infty}(\mathbb{R}^3)^2} \leq C \sigma^{(k-2)/3}, \quad k = 0, 1, \quad (3.17)$$

where \tilde{A}_j^\sharp is given by Lemma 3.1, and

$$\|A_{j,\sigma}^\sharp\|_{W^{k,\infty}(\mathbb{R}^3)^2} \leq C\sigma^{(k-2)/3}\|\tilde{A}_j^\sharp\|_{W^{2,\infty}(\mathbb{R}^3)} \leq C\sigma^{(k-2)/3}, \quad k \geq 2. \quad (3.18)$$

For further use, we notice from (3.4) and from (3.18) with $k = 2$, that the following estimate

$$\|b_{j,\sigma}\|_{W^{2,\infty}(\mathbb{R} \times \Omega)} + \|\partial_t b_{j,\sigma}\|_{W^{2,\infty}(\mathbb{R} \times \Omega)} \leq C, \quad j = 1, 2, \quad (3.19)$$

holds uniformly in $\sigma > 0$, for some constant $C > 0$ which is independent of σ . Moreover, it can be checked through direct calculation from (3.4), that

$$\begin{aligned} \theta \cdot \nabla_{x'} b_{j,\sigma}(t, x) &= -i \left(\sum_{m=1}^2 \theta_m \int_0^t \sum_{k=1}^2 \theta_k \partial_{x_k} a_{j,m,\sigma}(x' - s\theta, x_3) ds \right) b_j(t, x) \\ &= i \left(\sum_{k=1}^2 \theta_k \int_0^t \frac{d}{ds} a_{j,k,\sigma}(x' - s\theta, x_3) ds \right) b_j(t, x) \\ &= i \left(\theta \cdot A_{j,\sigma}^\sharp(x' - t\theta, x_3) - \theta \cdot A_{j,\sigma}^\sharp(x', x_3) \right) b_j(t, x), \end{aligned}$$

for all $(t, x) \in Q$, from where we see that $b_{j,\sigma}$ is solution to the transport equation

$$(\partial_t + \theta \cdot \nabla_{x'} + i\theta \cdot A_{j,\sigma}^\sharp) b_{j,\sigma} = 0, \quad \text{in } Q, \quad \sigma \in \mathbb{R}_+^*, \quad j = 1, 2. \quad (3.20)$$

We turn now to building suitable GO solutions to the magnetic Schrödinger equation of (1.1).

3.2. Building GO solutions to magnetic Schrödinger equations. For $j = 1, 2$, we seek GO solutions to the magnetic Schrödinger equation of (1.1) with (A, q) replaced by (A_j, q_j) , obeying (3.3)–(3.6), where the function $A_{j,\sigma}^\sharp$, appearing in (3.4), is the smooth magnetic potential described by Lemma 3.3. This requires that the functions Φ_j , appearing in 3.3, be preliminarily defined more explicitly. To do that, we set $B(0, r) := \{x' \in \mathbb{R}^2; |x'| < r\}$ for all $r > 0$, and take $R > 1$ so large that $\bar{\omega} \subset B(0, R - 1)$, where $\tilde{\omega}$ is the same as in Lemma 3.1. Next, we pick $\phi_j \in \mathcal{C}_0^\infty(\mathbb{R}^3)$, such that

$$\text{supp } \phi_j(\cdot, x_3) \subset \mathcal{D}_R := B(0, R + 1) \setminus \overline{B(0, R)}, \quad x_3 \in \mathbb{R}, \quad (3.21)$$

and put

$$\Phi_j(t, x) := \phi_j(x' - t\theta, x_3), \quad (t, x) \in \mathbb{R} \times \mathbb{R}^3. \quad (3.22)$$

It is apparent from (3.21) and the embedding $\omega \subset B(0, R - 1)$, that

$$\text{supp } \phi_j(\cdot, x_3) \cap \omega = \emptyset, \quad x_3 \in \mathbb{R}, \quad (3.23)$$

and from (3.22), that Φ is solution to the transport equation (3.5).

In the sequel, we choose $\sigma > \sigma_* := (R + 1)/T$, in such a way that

$$\text{supp } \Phi_j(\pm 2\sigma t, \cdot, x_3) \cap \omega = \text{supp } \phi_j(\cdot \mp 2\sigma t\theta, x_3) \cap \omega = \emptyset, \quad (t, x_3) \in [T, +\infty) \times \mathbb{R}. \quad (3.24)$$

Notice that upon possibly enlarging R , we may assume that $\sigma_* \geq 1$, which will always be the case in the remaining part of this text.

Next, for $k \in \mathbb{N}$, we introduce the following subspace of $H^k(\mathbb{R}^3)$,

$$\mathcal{H}_\theta^k := \{\phi \in H^k(\mathbb{R}^3); \theta \cdot \nabla_{x'} \cdot \phi \in H^k(\mathbb{R}^3) \text{ and } \text{supp } \phi(\cdot, x_3) \subset \mathcal{D}_R \text{ for a.e. } x_3 \in \mathbb{R}\},$$

endowed with the norm

$$N_{k,\theta}(\phi) := \|\phi\|_{H^k(\mathbb{R}^3)} + \|\theta \cdot \nabla_{x'} \phi\|_{H^k(\mathbb{R}^3)}, \quad \phi \in \mathcal{H}_\theta^2. \quad (3.25)$$

For notational simplicity, we put

$$\mathcal{N}_{\theta,\sigma}(\phi) := N_{2,\theta}(\phi) + \sigma^{1/3} N_{0,\theta}(\phi). \quad (3.26)$$

The coming statement claims existence of GO solutions $u_{j,\sigma}$, expressed by (3.3), with $L^2(0, T; H^k(\Omega))$ -norm of correction term $\psi_{j,\sigma}$ bounded by $\mathcal{N}_{\theta,\sigma}(\phi_j)/\sigma^{1-k}$ for $k = 0, 1$.

Proposition 3.1. *Let $M > 0$, and let $A_j \in W^{2,\infty}(\Omega, \mathbb{R}^3)$ and $q_j \in W^{1,\infty}(\Omega, \mathbb{R})$, $j = 1, 2$, satisfy (3.1)-(3.2). Then, for all $\sigma > \sigma_*$, there exists $u_{j,\sigma} \in C^1([0, T], L^2(\Omega)) \cap C([0, T], H^2(\Omega))$ obeying (3.3)-(3.6), where Φ_j is defined by (3.21)-(3.22), such that we have*

$$(i\partial_t + \Delta_{A_j} + q_j) u_{j,\sigma} = 0 \text{ in } Q,$$

and the correction term satisfies $\psi_{j,\sigma} = 0$ on Σ , for $j = 1, 2$, and $\psi_{1,\sigma}(T, \cdot) = \psi_{2,\sigma}(0, \cdot) = 0$ in Ω . Moreover, the following estimate

$$\sigma \|\psi_{j,\sigma}\|_{L^2(Q)} + \|\nabla \psi_{j,\sigma}\|_{L^2(Q)^3} \leq C \mathcal{N}_{\theta,\sigma}(\phi_j), \quad j = 1, 2, \quad (3.27)$$

holds for some constant $C > 0$ depending only on T , ω , and M , where the function $\phi_j \in C_0^\infty(\mathbb{R}^3)$ fulfills (3.21).

Proof. We prove the result for $j = 2$, the case $j = 1$ being obtained in the same way.

In light of (3.3)–(3.5) and the identity $(i\partial_t + \Delta_{A_2} + q_2)u_{2,\sigma} = 0$ imposed on $u_{2,\sigma}$ in Q , we seek a solution $\psi_{2,\sigma}$ to the following IBVP

$$\begin{cases} (i\partial_t + \Delta_{A_2} + q_2) \psi_{2,\sigma} = g_\sigma, & \text{in } Q, \\ \psi_{2,\sigma}(0, \cdot) = 0, & \text{in } \Omega, \\ \psi_{2,\sigma} = 0, & \text{on } \Sigma, \end{cases} \quad (3.28)$$

where

$$g_\sigma := -(i\partial_t + \Delta_{A_2} + q_2)(w_\sigma \varphi_\sigma),$$

with

$$w_\sigma(t, x') := e^{i\sigma(x' \cdot \theta - \sigma t)} \text{ and } \varphi_\sigma(t, x) := \vartheta_\sigma(2\sigma t, x), \text{ where } \vartheta_\sigma := \Phi_2 b_{2,\sigma}. \quad (3.29)$$

Next, taking into account that $(i\partial_t + \Delta_{A_2} + q_2)w_\sigma = (i\nabla \cdot A_2 - |A_2|^2 - 2\sigma\theta \cdot A_2^\sharp + q_2)w_\sigma$, and recalling from (3.5) and (3.20) that $i(\partial_t + 2\sigma\theta \cdot \nabla_{x'})\varphi_\sigma = 2\sigma\theta \cdot A_{2,\sigma}^\sharp \varphi_\sigma$, we get by straightforward computations that

$$g_\sigma(t, x) = -w_\sigma(t, x) \sum_{m=0,1} g_{m,\sigma}(2\sigma t, x), \text{ with } g_{0,\sigma} := (\Delta_{A_2} + q_2)\vartheta_\sigma, \quad g_{1,\sigma} := 2\sigma\theta \cdot (A_{2,\sigma}^\sharp - A_2^\sharp)\vartheta_\sigma. \quad (3.30)$$

As $g_\sigma \in W^{1,1}(0, T; L^2(\Omega))$, by (3.21)-(3.22), we know from Lemma 2.2 that (3.28) admits a unique solution $\psi_{2,\sigma} \in C^1([0, T], L^2(\Omega)) \cap C([0, T], H_0^1(\Omega) \cap H^2(\Omega))$. Moreover, since

$$\psi_{2,\sigma}(t, x) = -i \int_0^t e^{-i(t-s)\mathcal{H}_{A_2, q_2}} g_\sigma(s, x) ds, \quad (t, x) \in Q,$$

where \mathcal{H}_{A_2, q_2} is the self-adjoint operator acting in $L^2(\Omega)$, which is defined in Subsection 2.1, we have

$$\|\psi_{2,\sigma}(t, \cdot)\|_{L^2(\Omega)} \leq \int_0^t \|e^{-i(t-s)\mathcal{H}_{A_2, q_2}} g_\sigma(s, \cdot)\|_{L^2(\Omega)} ds \leq \|g_\sigma\|_{L^1(0, T; L^2(\Omega))},$$

uniformly in $t \in (0, T)$. This entails $\|\psi_{2,\sigma}\|_{L^2(Q)} \leq T^{1/2} \|g_\sigma\|_{L^1(0, T; L^2(\Omega))}$, which together with (3.30), yields

$$\|\psi_{2,\sigma}\|_{L^2(Q)} \leq T^{1/2} \sum_{m=0,1} \int_0^T \|g_{m,\sigma}(2\sigma t, \cdot)\|_{L^2(\Omega)} dt \leq \sigma^{-1} T^{1/2} \sum_{m=0,1} \|g_{m,\sigma}\|_{L^1(\mathbb{R}, L^2(\Omega))}. \quad (3.31)$$

We are left with the task of bounding each term $\|g_{m,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))}$, for $m = 0, 1$, separately. We start with $m = 0$, and obtain

$$\begin{aligned}\|g_{0,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} &= \int_{\mathbb{R}} \|(\Delta_{A_2} + q_2)(\Phi_2 b_{2,\sigma})(s, \cdot)\|_{L^2(\Omega)} ds \\ &\leq C \|b_{2,\sigma}\|_{W^{2,\infty}(\mathbb{R} \times \Omega)} \|\phi_2\|_{H^2(\mathbb{R}^3)} \leq C \|\phi_2\|_{H^2(\mathbb{R}^3)},\end{aligned}\quad (3.32)$$

by combining estimate (3.19) with definitions (3.21)-(3.22) and (3.30). Next, applying (3.17) with $k = 0$, we get that

$$\|g_{1,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} \leq C \sigma \|A_{2,\sigma}^\# - A_2^\#\|_{L^\infty(\Omega)} \|\phi_2\|_{L^2(\mathbb{R}^3)} \leq C \sigma^{1/3} \|\phi_2\|_{L^2(\mathbb{R}^3)}, \quad (3.33)$$

Putting (3.31)–(3.33) together, and recalling (3.25), we find that

$$\sigma \|\psi_{2,\sigma}\|_{L^2(Q)} \leq C \left(\|\phi_2\|_{H^2(\mathbb{R}^3)} + \sigma^{1/3} \|\phi_2\|_{L^2(\mathbb{R}^3)} \right) \leq C \left(N_{2,\theta}(\phi_2) + \sigma^{1/3} N_{0,\theta}(\phi_2) \right). \quad (3.34)$$

It remains to bound $\|\nabla \psi_{2,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))}$ from above. To do that, we apply [10, Lemma 3.2], which is permitted since $g_\sigma(0, \cdot) = 0$, with $\varepsilon = \sigma^{-1}$, getting

$$\begin{aligned}\|\nabla \psi_{2,\sigma}(t, \cdot)\|_{L^2(\Omega)} &\leq C \left(\sigma \|g_\sigma\|_{L^1(0,T;L^2(\Omega))} + \sigma^{-1} \|\partial_t g_\sigma\|_{L^1(0,T;L^2(\Omega))} \right) \\ &\leq C \sum_{m=0,1} \left(\sigma \int_0^T \|g_{m,\sigma}(2\sigma t, \cdot)\|_{L^2(\Omega)} dt + \int_0^T \|\partial_t g_{m,\sigma}(2\sigma t, \cdot)\|_{L^2(\Omega)} dt \right) \\ &\leq C \sum_{m=0,1} \left(\|g_{m,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} + \|\partial_t g_{m,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} \right),\end{aligned}\quad (3.35)$$

for every $t \in (0, T)$, according to (3.29)-(3.30). Further, as we have

$$\partial_t g_{0,\sigma}(t, x) = -(\Delta_{A_2} + q_2)\theta \cdot \left(\nabla_{x'} \phi_2 + i A_{2,\sigma}^\# \phi_2 \right) (x' - t\theta, x_3) b_{2,\sigma}(t, x),$$

for a.e. $(t, x) \in \mathbb{R} \times \Omega$, by direct computation, we obtain

$$\|\partial_t g_{0,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} \leq C N_{2,\theta}(\phi_2), \quad (3.36)$$

from (3.4), (3.18) with $k = 2$, (3.19) with $j = 2$, (3.21)-(3.22) and (3.29)-(3.30). Similarly, as

$$\partial_t g_{1,\sigma}(t, x) = -2\sigma\theta \cdot (A_2^\# - A_{2,\sigma}^\#)(x)\theta \cdot \left(\nabla_{x'} \phi_2 + i A_{2,\sigma}^\# \phi_2 \right) (x' - t\theta, x_3) b_{2,\sigma}(t, x),$$

for a.e. $(t, x) \in \mathbb{R} \times \Omega$, we find that

$$\|\partial_t g_{1,\sigma}\|_{L^1(\mathbb{R},L^2(\Omega))} \leq C \sigma \|A_2^\# - A_{2,\sigma}^\#\|_{L^\infty(\Omega)} N_{0,\theta}(\phi_2) \leq C \sigma^{1/3} N_{0,\theta}(\phi_2), \quad (3.37)$$

according to (3.17) with $j = 2$ and $k = 0$. Thus, we infer from (3.32)-(3.33) and (3.35)–(3.37), that

$$\|\nabla \psi_{2,\sigma}(t, \cdot)\|_{L^2(\Omega)^3} \leq C \left(N_{2,\theta}(\phi_2) + \sigma^{1/3} N_{0,\theta}(\phi_2) \right), \quad t \in (0, T), \quad \sigma > \sigma_*.$$

This and (3.34) yield (3.27) with $j = 2$, upon recalling the definition (3.26). \square

Let us now establish for further use that we may substitute $\sigma^{-1/6} u_{j,\sigma}$ for $\psi_{j,\sigma}$ in the estimate (3.27).

Corollary 3.4. *For $j = 1, 2$, let q_j , A_j , ϕ_j , and $u_{j,\sigma}$, be the same as in Proposition 3.1. Then, there exists a constant $C > 0$, depending only on T , ω and M , such that the estimate*

$$\sigma \|u_{j,\sigma}\|_{L^2(Q)} + \|\nabla u_{j,\sigma}\|_{L^2(Q)^3} \leq C \sigma^{1/6} \mathcal{N}_{\theta,\sigma}(\phi_j), \quad j = 1, 2, \quad (3.38)$$

holds for all $\sigma > \sigma_*$.

Proof. Notice from (3.22) and (3.24) that

$$\int_0^T \|\Phi_j(2\sigma t, \cdot)\|_{H^k(\Omega)}^2 dt = \int_0^{+\infty} \|\Phi_j(2\sigma t, \cdot)\|_{H^k(\Omega)}^2 dt = (2\sigma)^{-1} \int_0^{2R} \|\Phi_j(s, \cdot)\|_{H^k(\Omega)}^2 ds,$$

so we have

$$\|\Phi_j(2\sigma \cdot, \cdot)\|_{L^2(0,T;H^k(\Omega))} \leq R^{1/2} \sigma^{-1/2} \|\phi_j\|_{H^k(\mathbb{R}^3)}, \quad j = 1, 2, \quad k \in \mathbb{N}. \quad (3.39)$$

From this, (3.3), (3.19) and (3.25)–(3.27), it follows for each $j = 1, 2$, that

$$\begin{aligned} \|u_{j,\sigma}\|_{L^2(Q)} &\leq \|b_{j,\sigma}\|_{L^\infty(\mathbb{R} \times \Omega)} \|\Phi_j(2\sigma \cdot, \cdot)\|_{L^2(Q)} + \|\psi_{j,\sigma}\|_{L^2(Q)} \\ &\leq C \left(\sigma^{-1/2} \|\phi_j\|_{L^2(\mathbb{R}^3)} + \sigma^{-1} \mathcal{N}_{\theta,\sigma}(\phi_j) \right) \leq C \sigma^{-5/6} \mathcal{N}_{\theta,\sigma}(\phi_j), \end{aligned}$$

and

$$\begin{aligned} &\|\nabla u_{j,\sigma}\|_{L^2(Q)^3} \\ &\leq \|b_{j,\sigma}\|_{W^{1,\infty}(\mathbb{R} \times \Omega)} \left(\sigma \|\Phi_j(2\sigma \cdot, \cdot)\|_{L^2(Q)} + \|\Phi_j(2\sigma \cdot, \cdot)\|_{L^2(0,T;H^1(\Omega))} \right) + \|\nabla \psi_{j,\sigma}\|_{L^2(Q)^3} \\ &\leq C \left(\sigma^{1/2} \|\phi_j\|_{L^2(\mathbb{R}^3)} + \sigma^{-1/2} \|\phi_j\|_{H^1(\mathbb{R}^3)} + \mathcal{N}_{\theta,\sigma}(\phi_j) \right) \leq C \sigma^{1/6} \mathcal{N}_{\theta,\sigma}(\phi_j), \end{aligned}$$

which yields (3.38). \square

In the coming subsection we probe the medium with the GO solutions described in Proposition 3.1 in order to upper bound the transverse magnetic potential in terms of suitable norm of the DN map.

3.3. Probing the medium with GO solutions. Let us introduce

$$\tilde{A}^\sharp := \tilde{A}_2^\sharp - \tilde{A}_1^\sharp, \text{ and } A_\sigma^\sharp := A_{2,\sigma}^\sharp - A_{1,\sigma}^\sharp, \quad \sigma > 0, \quad (3.40)$$

where the functions \tilde{A}_j^\sharp and $A_{j,\sigma}^\sharp$, $j = 1, 2$, are defined in Lemma 3.1 and Lemma 3.3, respectively. Evidently, \tilde{A}^\sharp is the function $A_2^\sharp - A_1^\sharp$, extended by zero outside Ω , and we have

$$\|A_\sigma^\sharp - \tilde{A}^\sharp\|_{W^{1,\infty}(\mathbb{R}^3)^2} \leq \sum_{j=1,2} \|A_{j,\sigma}^\sharp - \tilde{A}_j^\sharp\|_{W^{1,\infty}(\mathbb{R}^3)^2} \leq 2C\sigma^{-1/3}, \quad \sigma > 0, \quad (3.41)$$

from (3.17) with $k = 1$. Thus, writing $A_\sigma^\sharp = (a_{1,\sigma}, a_{2,\sigma})$ and $\tilde{A}^\sharp = (\tilde{a}_1, \tilde{a}_2)$, it follows readily from (3.16) that

$$a_{i,\sigma}(x) = \int_{\mathbb{R}^3} \chi_\sigma(x-y) (\tilde{a}_i(y) + (x-y) \cdot \nabla \tilde{a}_i(y)) dy, \quad x \in \mathbb{R}^3, \quad i = 1, 2. \quad (3.42)$$

The main purpose of this subsection is the following technical result.

Lemma 3.5. *Let $M > 0$ and $\theta \in \mathbb{S}^1$ be fixed. For $j = 1, 2$, let $A_j \in W^{2,\infty}(\Omega, \mathbb{R})^3$, $q_j \in W^{1,\infty}(\Omega, \mathbb{R}^3)$, obey (3.1)–(3.2), and let ϕ_j be defined by (3.21). Then, for every $\sigma > \sigma_*$, there exists a constant $C > 0$, depending only on T , ω , and M , such that we have*

$$\begin{aligned} &\sigma \left| \int_{(0,T) \times \mathbb{R}^3} \theta \cdot \tilde{A}^\sharp(x) (\overline{\phi_1} \phi_2)(x' - 2\sigma t \theta, x_3) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x) dx' dx_3 dt \right| \\ &\leq C \left(\sigma^5 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-5/6} \right) \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2), \end{aligned} \quad (3.43)$$

where $\|\cdot\|$ stands for the usual norm in $\mathcal{B}(H^{2,1}(\Sigma), L^2(\Sigma))$, and \tilde{A}^\sharp is given by (3.40).

Proof. We proceed in two steps. The first step is to establish a suitable orthogonality identity for $A := A_2 - A_1$ and $V := i\nabla \cdot A - (|A_2|^2 - |A_1|^2) + q_2 - q_1$, which is the key ingredient in the derivation of the estimate (3.43), presented in the second step.

Step 1: Orthogonality identity. We probe the system with the GO functions $u_{j,\sigma}$, $j = 1, 2$, given by Proposition 3.1, and recall for further use that $u_{j,\sigma} \in \mathcal{C}^1([0, T], L^2(\Omega)) \cap \mathcal{C}([0, T], H^2(\Omega))$ is expressed by (3.3) and satisfies the following equation

$$(i\partial_t + \Delta_{A_j} + q_j) u_{j,\sigma} = 0, \text{ in } Q. \quad (3.44)$$

Since $A_{2,\sigma}^\# \in W^{2,\infty}(\Omega)^2$ and $\phi_2 \in C_0^\infty(\mathbb{R}^3)$, it follows readily from (3.3)-(3.4) and (3.22) that $u_{2,\sigma} - \psi_{2,\sigma} \in C^\infty([0, T], W^{2,\infty}(\Omega))$. Thus, $F := -(i\partial_t + \Delta_{A_1} + q_1)(u_{2,\sigma} - \psi_{2,\sigma}) \in W^{1,1}(0, T; L^2(\Omega))$, and there is consequently a unique solution $z \in \mathcal{C}^1([0, T], L^2(\Omega)) \cap \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega))$ to the IBVP

$$\begin{cases} (i\partial_t + \Delta_{A_1} + q_1) z = F, & \text{in } Q, \\ z(0, \cdot) = 0, & \text{in } \Omega, \\ z = 0, & \text{on } \Sigma, \end{cases} \quad (3.45)$$

in virtue of Lemma 2.2. Further, as $(u_{2,\sigma} - \psi_{2,\sigma})(0, \cdot) = 0$ in Ω , by (3.22)-(3.23), we infer from (3.45) that $v := z + u_{2,\sigma} - \psi_{2,\sigma} \in \mathcal{C}^1([0, T], L^2(\Omega)) \cap \mathcal{C}([0, T], H^2(\Omega))$ verifies

$$\begin{cases} (i\partial_t + \Delta_{A_1} + q_1) v = 0, & \text{in } Q, \\ v(0, \cdot) = 0, & \text{in } \Omega, \\ v = f_\sigma, & \text{on } \Sigma, \end{cases} \quad (3.46)$$

where we have set

$$f_\sigma(t, x) := u_{2,\sigma}(t, x) - \psi_{2,\sigma}(t, x) = (\Phi_2 b_{2,\sigma})(2\sigma t, x) e^{i\sigma(x' \cdot \theta - \sigma t)}, \quad (t, x) \in \Sigma. \quad (3.47)$$

From this and Proposition 3.1, it then follows that $w := v - u_{2,\sigma}$ is the $\mathcal{C}^1([0, T], L^2(\Omega)) \cap \mathcal{C}([0, T], H_0^1(\Omega) \cap H^2(\Omega))$ -solution to the IBVP

$$\begin{cases} (i\partial_t + \Delta_{A_1} + q_1) w = 2iA \cdot \nabla u_{2,\sigma} + V u_{2,\sigma}, & \text{in } Q, \\ w(0, \cdot) = 0, & \text{in } \Omega, \\ w = 0, & \text{on } \Sigma, \end{cases} \quad (3.48)$$

In light of (3.48), we deduce from (3.44) with $j = 1$, upon applying the Green formula, that

$$\begin{aligned} \langle 2iA \cdot \nabla u_{2,\sigma} + V u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} &= \langle (i\partial_t + \Delta_{A_1} + q_1) w, u_{1,\sigma} \rangle_{L^2(Q)} \\ &= \langle (\partial_\nu + iA_1 \cdot \nu) w, u_{1,\sigma} \rangle_{L^2(\Sigma)}. \end{aligned} \quad (3.49)$$

Next, taking into account that $A_1 = A_2$ on $\partial\Omega$, from (3.2), we see that

$$\begin{aligned} (\partial_\nu + iA_1 \cdot \nu) w &= (\partial_\nu + iA_1 \cdot \nu) v - (\partial_\nu + iA_1 \cdot \nu) u_{2,\sigma} \\ &= (\partial_\nu + iA_1 \cdot \nu) v - (\partial_\nu + iA_2 \cdot \nu) u_{2,\sigma} \\ &= (\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}) f_\sigma, \end{aligned}$$

according to (3.47) and the last line of (3.46). This and (3.49) yield the following orthogonality identity

$$2i \langle A \cdot \nabla u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} + \langle V u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} = \langle (\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}) f_\sigma, g_\sigma \rangle_{L^2(\Sigma)}, \quad (3.50)$$

with

$$g_\sigma(t, x) := u_{1,\sigma}(t, x) = u_{1,\sigma}(t, x) - \psi_{1,\sigma}(t, x) = (\Phi_1 b_{1,\sigma})(2\sigma t, x) e^{i\sigma(x' \cdot \theta - 2\sigma t)}, \quad (t, x) \in \Sigma. \quad (3.51)$$

Having established (3.50), we turn now to proving the estimate (3.43).

Step 2: Derivation of (3.43). In light of (3.3), we have

$$\langle A \cdot \nabla u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} = I_\sigma + i\sigma \int_Q \theta \cdot A^\sharp(x) (\overline{\Phi_1 \Phi_2})(2\sigma t, x) (\overline{b_{1,\sigma} b_{2,\sigma}})(2\sigma t, x) dx dt, \quad (3.52)$$

with

$$\begin{aligned} I_\sigma &:= \int_Q A \cdot \nabla (\Phi_2 b_{2,\sigma})(2\sigma t, x) \left((\overline{\Phi_1 b_{1,\sigma}})(2\sigma t, x) + e^{i\sigma(x' \cdot \theta - \sigma t)} \overline{\psi_{1,\sigma}}(t, x) \right) dx dt \\ &\quad + \int_Q A \cdot \nabla \psi_{2,\sigma}(t, x) \left(e^{i\sigma(x' \cdot \theta - \sigma t)} (\overline{\Phi_1 b_{1,\sigma}})(2\sigma t, x) + \overline{\psi_{1,\sigma}}(t, x) \right) dx dt \\ &\quad + i\sigma \int_Q \theta \cdot A^\sharp(x) (\Phi_2 b_{2,\sigma})(2\sigma t, x) \overline{\psi_{1,\sigma}}(t, x) e^{i\sigma(x' \cdot \theta - \sigma t)} dx dt. \end{aligned}$$

We infer from (3.19), (3.27), and (3.39), that

$$|I_\sigma| \leq C \sigma^{-5/6} \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2), \quad \sigma > \sigma_*.$$

Putting this together with (3.50) and (3.52), we find that

$$\begin{aligned} &\sigma \left| \int_Q \theta \cdot A^\sharp(x) (\phi_2 \overline{\phi_1})(x' - 2\sigma t \theta, x_3) (\overline{b_{2,\sigma} b_{1,\sigma}})(2\sigma t, x) dx' dx_3 dt \right| \\ &\leq C \left(|\langle V u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)}| + |\langle (\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}) f_\sigma, g_\sigma \rangle_{L^2(\Sigma)}| + \sigma^{-5/6} \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2) \right) \end{aligned} \quad (3.53)$$

Next, we notice from (3.38) that

$$|\langle V u_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)}| \leq C \sigma^{-5/3} \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2). \quad (3.54)$$

Moreover, in view of (3.47) and (3.51), we have

$$\begin{aligned} |\langle (\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}) f_\sigma, g_\sigma \rangle_{L^2(\Sigma)}| &\leq \|\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}\| \|f_\sigma\|_{H^{2,1}(\Sigma)} \|g_\sigma\|_{L^2(\Sigma)} \\ &\leq \|\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}\| \|u_{2,\sigma} - \psi_{2,\sigma}\|_{H^{2,1}(\Sigma)} \|u_{1,\sigma} - \psi_{1,\sigma}\|_{L^2(\Sigma)}, \end{aligned}$$

with

$$\begin{aligned} \|u_{1,\sigma} - \psi_{1,\sigma}\|_{L^2(\Sigma)} &\leq \|u_{1,\sigma} - \psi_{1,\sigma}\|_{L^2(0,T;H^1(\Omega))} \\ &\leq C \sigma \|\Phi_1(2\sigma \cdot, \cdot)\|_{L^2(0,T;H^1(\Omega))} \|b_{1,\sigma}\|_{W^{1,\infty}(\mathbb{R} \times \Omega)} \\ &\leq C \sigma^{1/2} \mathcal{N}_{\theta,\sigma}(\phi_1), \end{aligned}$$

and

$$\begin{aligned} \|u_{2,\sigma} - \psi_{2,\sigma}\|_{H^{2,1}(\Sigma)} &\leq C (\|u_{2,\sigma} - \psi_{2,\sigma}\|_{H^2(0,T;H^1(\Omega))} + \|u_{2,\sigma} - \psi_{2,\sigma}\|_{L^2(0,T;H^2(\Omega))}) \\ &\leq C \sigma^5 \|\Phi_2(2\sigma \cdot, \cdot)\|_{L^2(0,T;H^2(\Omega))} (\|b_{2,\sigma}\|_{W^{2,\infty}(\mathbb{R} \times \Omega)} + \|\partial_t b_{2,\sigma}\|_{W^{2,\infty}(\mathbb{R} \times \Omega)}) \\ &\leq C \sigma^{9/2} \mathcal{N}_{\theta,\sigma}(\phi_2), \end{aligned}$$

according to (3.3), (3.19), (3.25), and (3.39). As a consequence, we have

$$|\langle (\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}) f_\sigma, g_\sigma \rangle_{L^2(\Sigma)}| \leq C \sigma^5 \|\Lambda_{A_1,q_1} - \Lambda_{A_2,q_2}\| \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2). \quad (3.55)$$

This and (3.53)-(3.54) yield (3.43). \square

4. PRELIMINARY ESTIMATES

4.1. **X-ray transform.** In this subsection we estimate the partial X-ray transform in \mathbb{R}^3 , of the functions

$$\tilde{\rho}_j(x', x_3) := \theta \cdot \frac{\partial \tilde{A}^\sharp}{\partial x_j}(x) = \sum_{i=1,2} \theta_i \frac{\partial \tilde{a}_i}{\partial x_j}(x), \quad x \in \mathbb{R}^3, \quad j = 1, 2, 3, \quad (4.1)$$

in terms of the DN map. We recall that the partial X-ray transform in the direction $\theta \in \mathbb{S}^1$, of a function

$$f \in \mathcal{X} := \{\varphi \in L^1_{\text{loc}}(\mathbb{R}^3); \quad x' \mapsto \varphi(x', x_3) \in L^1(\mathbb{R}^2) \text{ for a.e. } x_3 \in \mathbb{R}\}, \quad (4.2)$$

is defined as

$$\mathcal{P}(f)(\theta, x', x_3) := \int_{\mathbb{R}} f(x' + s\theta, x_3) ds, \quad x' \in \mathbb{R}^2, \quad x_3 \in \mathbb{R}. \quad (4.3)$$

The X-ray transform stability estimate is as follows.

Lemma 4.1. *Let $M > 0$, and let A_j and q_j , for $j = 1, 2$, be as in Proposition 3.1. Then, there exists a constant $C > 0$, depending only on T , ω and M , such that for all $\theta \in \mathbb{S}^1$, all $\xi' \in \mathbb{R}^2$, and all $\phi \in C_0^\infty(\mathbb{R}^3)$ satisfying $\text{supp } \phi(\cdot, x_3) \subset \mathcal{D}_R^-(\theta) := \{x' \in \mathcal{D}_R, \quad x' \cdot \theta \leq 0\}$ for every $x_3 \in \mathbb{R}$, the estimate*

$$\begin{aligned} & \left| \int_{\mathbb{R}^3} \phi^2(x) \mathcal{P}(\tilde{\rho}_j)(\theta, x', x_3) \exp\left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds\right) dx \right| \\ & \leq C \left(\sigma^5 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-5/6} \right) \mathcal{N}_{\theta, \sigma}(\phi) \mathcal{N}_{\theta, \sigma}(\partial_{x_j} \phi), \end{aligned} \quad (4.4)$$

holds uniformly in $\sigma > \sigma_*$ and $j = 1, 2, 3$.

Proof. Let $\phi_j \in C_0^\infty(\mathbb{R}^3)$, $j = 1, 2$, be supported in $\mathcal{D}_R \times \mathbb{R}$. Bearing in mind that $\bar{\omega} \subset B(0, R-1)$, we infer from (3.42) that \tilde{A}^\sharp and A_σ^\sharp are both supported in $B(0, R) \times \mathbb{R}$. Further, as $|x' - 2\sigma t\theta| > 2\sigma_* T - R > R+1$ for all $x' \in B(0, R)$ and $t > T$, we see that

$$\tilde{A}^\sharp(x)(\bar{\phi}_1 \phi_2)(x' - 2\sigma t\theta, x_3) = A_\sigma^\sharp(x)(\bar{\phi}_1 \phi_2)(x' - 2\sigma t\theta, x_3) = 0, \quad x = (x', x_3) \in \mathbb{R}^3, \quad t > T,$$

As a consequence we have

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^3} \theta \cdot \left(\tilde{A}^\sharp(x) - A_\sigma^\sharp(x) \right) (\bar{\phi}_1 \phi_2)(x' - 2\sigma t\theta, x_3) (\bar{b}_{1, \sigma} b_{2, \sigma})(2\sigma t, x) dx dt \\ & = \int_0^{+\infty} \int_{\mathbb{R}^3} \theta \cdot \left(\tilde{A}^\sharp(x) - A_\sigma^\sharp(x) \right) (\bar{\phi}_1 \phi_2)(x' - 2\sigma t\theta, x_3) (\bar{b}_{1, \sigma} b_{2, \sigma})(2\sigma t, x) dx dt. \end{aligned}$$

Next, making the substitution $s = \sigma t$ in the above integral, we get that

$$\begin{aligned} & \sigma \left| \int_0^T \int_{\mathbb{R}^3} \theta \cdot \left(\tilde{A}^\sharp(x) - A_\sigma^\sharp(x) \right) (\bar{\phi}_1 \phi_2)(x' - 2\sigma t\theta, x_3) (\bar{b}_{1, \sigma} b_{2, \sigma})(2\sigma t, x) dx dt \right| \\ & = \left| \int_0^{+\infty} \int_{B(0, R) \times \mathbb{R}} \theta \cdot \left(\tilde{A}^\sharp(x) - A_\sigma^\sharp(x) \right) (\bar{\phi}_1 \phi_2)(x' - 2s\theta, x_3) (\bar{b}_{1, \sigma} b_{2, \sigma})(2s, x) dx ds \right| \\ & \leq \|\tilde{A}^\sharp - A_\sigma^\sharp\|_{L^\infty(\mathbb{R}^3)^2} \int_0^{R+1} \int_{B(0, R) \times \mathbb{R}} |(\bar{\phi}_1 \phi_2)(x' - 2s\theta, x_3)| dx ds \\ & \leq \|\tilde{A}^\sharp - A_\sigma^\sharp\|_{L^\infty(\mathbb{R}^3)^2} \int_0^{R+1} \int_{\mathbb{R}^3} |(\bar{\phi}_1 \phi_2)(x' - 2s\theta, x_3)| dx ds \\ & \leq (R+1) \|\tilde{A}^\sharp - A_\sigma^\sharp\|_{L^\infty(\mathbb{R}^3)^2} \|\phi_1\|_{L^2(\mathbb{R}^3)} \|\phi_2\|_{L^2(\mathbb{R}^3)}. \end{aligned}$$

From this and (3.17) with $k = 0$, and (3.25), it follows that

$$\begin{aligned} & \left| \sigma \int_0^T \int_{\mathbb{R}^3} \theta \cdot \left(\tilde{A}^\sharp(x) - A_\sigma^\sharp(x) \right) (\overline{\phi_1} \phi_2)(x' - 2\sigma t \theta, x_3) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x) dx dt \right| \\ & \leq C \sigma^{-2/3} \|\phi_1\|_{L^2(\mathbb{R}^3)} \|\phi_2\|_{L^2(\mathbb{R}^3)} \leq C \sigma^{-4/3} \mathcal{N}_{\theta,\sigma}(\phi_1) \mathcal{N}_{\theta,\sigma}(\phi_2). \end{aligned} \quad (4.5)$$

On the other hand, since

$$\begin{aligned} (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x' + 2\sigma t \theta, x_3) &= \exp \left(-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(x' + (2\sigma t - s)\theta, x_3) ds \right) \\ &= \exp \left(-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right), \end{aligned}$$

for a.e. $(t, x) \in (0, T) \times \mathbb{R}^3$, we have

$$\begin{aligned} & \sigma \int_0^T \int_{\mathbb{R}^3} \theta \cdot A_\sigma^\sharp(x) (\overline{\phi_1} \phi_2)(x' - 2\sigma t \theta, x_3) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x) dx' dx_3 dt \\ &= \sigma \int_0^T \int_{\mathbb{R}^3} \theta \cdot A_\sigma^\sharp(x' + 2\sigma t \theta, x_3) (\overline{\phi_1} \phi_2)(x) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x' + 2\sigma t \theta, x_3) dx' dx_3 dt \\ &= \int_{\mathbb{R}^3} (\overline{\phi_1} \phi_2)(x) \left(\int_0^T \sigma \theta \cdot A_\sigma^\sharp(x' + 2\sigma t \theta, x_3) \exp \left(-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dt \right) dx' dx_3 \\ &= \frac{i}{2} \int_{\mathbb{R}^3} (\overline{\phi_1} \phi_2)(x) \left(\int_0^T \frac{d}{dt} \exp \left(-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dt \right) dx' dx_3 \\ &= \frac{i}{2} \int_{\mathbb{R}^3} (\overline{\phi_1} \phi_2)(x) \left(\exp \left(-i \int_0^{2\sigma T} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) - 1 \right) dx' dx_3. \end{aligned} \quad (4.6)$$

As A_σ^\sharp is supported in $B(0, R) \times \mathbb{R}$ and $|x' + s\theta| > 2\sigma_* T - (R + 1) > R$, for all $x' \in \mathcal{D}_R$ and all $s > 2\sigma T$, we have

$$\int_0^{2\sigma T} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds = \int_0^{+\infty} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds, \quad x' \in \mathcal{D}_R, \quad x_3 \in \mathbb{R}. \quad (4.7)$$

Similarly, as $|x' + s\theta|^2 = |x'|^2 + s^2 + 2sx' \cdot \theta > R^2$, for every $x' \in \mathcal{D}_R^-(\theta)$ and $s < 0$, we have

$$\int_{-\infty}^0 \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds = 0, \quad x' \in \mathcal{D}_R^-(\theta), \quad x_3 \in \mathbb{R}.$$

This and (4.7) entail

$$\int_0^{2\sigma T} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds = \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds, \quad x' \in \mathcal{D}_R^-(\theta), \quad x_3 \in \mathbb{R}. \quad (4.8)$$

Having seen this, we take $\phi_1 := \partial_{x_j} \overline{\phi}$, for $j = 1, 2, 3$, and $\phi_2 := \phi$, in (4.6), and find

$$\begin{aligned} & \sigma \int_0^T \int_{\mathbb{R}^3} \theta \cdot A_\sigma^\sharp(x) (\overline{\phi_1} \phi_2)(x' - 2\sigma t \theta, x_3) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x) dx' dx_3 dt \\ &= \frac{i}{4} \int_{\mathbb{R}^3} \partial_{x_j} \phi^2(x) \left(\exp \left(-i \int_0^{2\sigma T} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) - 1 \right) dx' dx_3 \\ &= -\frac{1}{4} \int_{\mathbb{R}^3} \phi^2(x) \left(\int_0^{2\sigma T} \theta \cdot \partial_{x_j} A_\sigma^\sharp(x' + s\theta, x_3) ds \right) \exp \left(-i \int_0^{2\sigma T} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dx \end{aligned} \quad (4.9)$$

upon integrating by parts. Taking into account that ϕ is supported in $\mathcal{D}_R^-(\theta) \times \mathbb{R}$, we deduce from (4.8) and (4.9), that

$$\begin{aligned} & \sigma \int_0^T \int_{\mathbb{R}^3} \theta \cdot A_\sigma^\sharp(x) (\overline{\phi_1} \phi_2)(x' - 2\sigma t \theta, x_3) (\overline{b_{1,\sigma}} b_{2,\sigma})(2\sigma t, x) dx' dx_3 dt \\ &= -\frac{1}{4} \int_{\mathbb{R}^3} \phi^2(x) \left(\int_{\mathbb{R}} \theta \cdot \partial_{x_j} A_\sigma^\sharp(x' + s\theta, x_3) ds \right) \exp \left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dx' dx_3 \\ &= -\frac{1}{4} \int_{\mathbb{R}^3} \phi^2(x) \mathcal{P}(\rho_{j,\sigma})(\theta, x', x_3) \exp \left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dx' dx_3. \end{aligned} \quad (4.10)$$

Here we used (4.3) and the notation

$$\rho_{j,\sigma}(x) := \theta \cdot \partial_{x_j} A_\sigma^\sharp(x) = \sum_{i=1,2} \theta_i \partial_{x_j} a_{i,\sigma}(x), \quad x \in \mathbb{R}^3, \quad j = 1, 2, 3.$$

Finally, using once more that the functions A_σ^\sharp and \tilde{A}^\sharp are supported in $B(0, R)$, we infer from (3.41) and (4.1)-(4.3), that

$$|(\mathcal{P}(\rho_{j,\sigma}) - \mathcal{P}(\tilde{\rho}_j))(\theta, x', x_3)| \leq C\sigma^{-1/3}, \quad (x', x_3) \in B(0, R) \times \mathbb{R},$$

for some positive constant C , depending only on ω and M . This entails that

$$\left| \int_{\mathbb{R}^3} \phi^2(x) (\mathcal{P}(\rho_{j,\sigma}) - \mathcal{P}(\tilde{\rho}_j))(\theta, x', x_3) \exp \left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, x_3) ds \right) dx' dx_3 \right| \leq C\sigma^{-1/3} \|\phi\|_{L^2(\mathbb{R}^3)}^2,$$

which, together with (3.43), (4.5), and (4.10), yields (4.4). \square

As will be seen in the coming section, the result of Lemma 4.1 is a key ingredient in the estimation of the partial Fourier transform of the aligned magnetic field, in terms of the DN map. To this purpose, we recall for all $f \in \mathcal{X}$, where \mathcal{X} is defined in (4.2), that the partial Fourier transform with respect to $x' \in \mathbb{R}^2$ of f , expresses as

$$\widehat{f}(\xi', x_3) := (2\pi)^{-1} \int_{\mathbb{R}^2} f(x', x_3) e^{-ix' \cdot \xi'} dx', \quad \xi' \in \mathbb{R}^2, \quad x_3 \in \mathbb{R}. \quad (4.11)$$

Further, setting $\theta^\perp := \{x' \in \mathbb{R}^2; x' \cdot \theta = 0\}$, we recall for further use from [9, Lemma 6.1], that $x' \mapsto \mathcal{P}(f)(\theta, x', x_3) \in L^1(\theta^\perp)$ for a.e. $x_3 \in \mathbb{R}$, and that

$$\widehat{\mathcal{P}(f)}(\theta, \xi', x_3) := (2\pi)^{-1/2} \int_{\theta^\perp} \mathcal{P}(f)(\theta, x', x_3) e^{-ix' \cdot \xi'} dx' = (2\pi)^{1/2} \widehat{f}(\xi', x_3), \quad \xi' \in \theta^\perp, \quad x_3 \in \mathbb{R}. \quad (4.12)$$

4.2. Aligned magnetic field estimation. Let us now estimate the Fourier transform of the aligned magnetic field

$$\tilde{\beta}(x) := (\partial_{x_1} \tilde{a}_2 - \partial_{x_2} \tilde{a}_1)(x), \quad x \in \mathbb{R}^3, \quad (4.13)$$

with the aid of Lemma 4.1. More precisely, we aim to establish the following result.

Lemma 4.2. *Let $M > 0$, and let A_j and q_j , for $j = 1, 2$, be as in Proposition 3.1. Then, there exist two constants $\epsilon \in (0, 1)$ and $C > 0$, both of them depending only on T , ω , and M , such that the estimates*

$$\|\widehat{\beta}(\xi', \cdot)\|_{L^\infty(\mathbb{R})} \leq C \langle \xi' \rangle^7 (\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-\epsilon}), \quad (4.14)$$

and

$$\|\partial_{x_3} \widehat{\beta}(\xi', \cdot)\|_{L^\infty(\mathbb{R})} \leq C \langle \xi' \rangle^8 (\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-\epsilon}), \quad (4.15)$$

hold for all $\sigma > \sigma_*$ and all $\xi' \in \mathbb{R}^2$, with $\langle \xi' \rangle := (1 + |\xi'|^2)^{1/2}$.

Proof. We shall only prove (4.14), the derivation of (4.15) being obtained in a similar fashion.

Fix $\theta \in \mathbb{S}^1 \cap \xi'^\perp$. We first introduce the following partition of $B(0, R) \cap \theta^\perp$. For $N \in \mathbb{N}^* := \{1, 2, \dots\}$ fixed, we pick x'_1, \dots, x'_N in $B(0, R + 1/2) \cap \theta^\perp$, and choose $\varphi_1, \dots, \varphi_N$ in $\mathcal{C}_0^\infty(\mathbb{R}^2, [0, 1])$, such that

$$\text{supp } \varphi_k \subset B(x'_k, 1/8) \cap \theta^\perp \text{ for } k = 1, \dots, N, \text{ and } \sum_{k=1}^N \varphi_k(x') = 1 \text{ for } x' \in B(0, R) \cap \theta^\perp. \quad (4.16)$$

Next, we set $r_{x'_k} := \left((R + 3/4)^2 - |x'_k|^2 \right)^{1/2}$, in such a way that

$$B(x'_k - r_{x'_k} \theta, 1/4) \subset \mathcal{D}_R^-(\theta), \quad k = 1, \dots, N. \quad (4.17)$$

In order to define a suitable set of test functions $\phi_{*,k}$, $k = 1, \dots, N$, we fix $x_3 \in \mathbb{R}$, pick a function $\alpha \in \mathcal{C}_0^\infty(\mathbb{R}, \mathbb{R}_+)$ which is supported in $(-1, 1)$ and normalized in $L^2(\mathbb{R})$, and put

$$\alpha_\sigma(s) := \sigma^\mu \alpha(\sigma^{2\mu}(x_3 - s)), \quad s \in \mathbb{R}, \quad (4.18)$$

for some positive real parameter μ , we shall make precise below. Then, the test function $\phi_{*,k}$ is defined for all $y = (y', y_3) \in \mathbb{R}^3$, by

$$\phi_{*,k}(y) := h(y' \cdot \theta + r_{x'_k}) e^{-\frac{i}{2} y' \cdot \xi'} \varphi_k^{1/2}(y' - (y' \cdot \theta) \theta) \exp\left(\frac{i}{2} \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(y' + s\theta, y_3) ds\right) \alpha_\sigma(y_3), \quad (4.19)$$

where $h \in \mathcal{C}_0^\infty(\mathbb{R})$ is supported in $(0, 1/8)$, and normalized in $L^2(\mathbb{R})$.

For every $y' \in \mathbb{R}^2 \setminus B(x'_k - r_{x'_k} \theta, 1/4)$, it is easily seen from the basic inequality

$$|y' - (x'_k - r_{x'_k} \theta)| \leq |y' - (y' \cdot \theta) \theta - x'_k| + |y' \cdot \theta + r_{x'_k}|,$$

that either of the two real numbers $|y' - (y' \cdot \theta) \theta - x'_k|$ or $|y' \cdot \theta + r_{x'_k}|$ is greater than $1/8$, and hence that $h(y' \cdot \theta + r_{x'_k}) \varphi_k^{1/2}(y' - (y' \cdot \theta) \theta) = 0$. As a consequence, we have

$$\text{supp } \phi_{*,k}(\cdot, y_3) \subset B(x'_k - r_{x'_k} \theta, 1/4) \subset \mathcal{D}_R^-(\theta), \quad y_3 \in \mathbb{R}, \quad k = 1, \dots, N, \quad (4.20)$$

directly from (4.17) and (4.19). Moreover, since

$$\theta \cdot \nabla_{y'} \left(\int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(y' + s\theta, y_3) ds \right) = \theta \cdot \int_{\mathbb{R}} \frac{d}{ds} A_\sigma^\sharp(y' + s\theta, y_3) ds = 0, \quad (y', y_3) \in \mathbb{R}^3,$$

we derive from Lemma 3.3 for all $m \in \mathbb{N}$, that

$$\langle \xi' \rangle \|\phi_{*,k}\|_{H^m(\mathbb{R}^3)} + \|\theta \cdot \nabla_{x'} \phi_{*,k}\|_{H^m(\mathbb{R}^3)} \leq C \langle \xi' \rangle^{m+1} \sigma^{2\mu m + \max(0, (m-2)/3)},$$

where C is a positive constant, independent of σ . Therefore, we have $N_{0,\theta}(\phi_{*,k}) \leq C \langle \xi' \rangle$ and $N_{2,\theta}(\phi_{*,k}) \leq C \langle \xi' \rangle^3 \sigma^{4\mu}$, whence

$$\mathcal{N}_{\theta,\sigma}(\phi_{*,k}) \leq C \langle \xi' \rangle^3 \sigma^{4\mu+1/3}. \quad (4.21)$$

Similarly, we find that

$$\langle \xi' \rangle \|\partial_{x_j} \phi_{*,k}\|_{H^m(\mathbb{R}^3)} + \|\theta \cdot \nabla_{x'} \partial_{x_j} \phi_{*,k}\|_{H^m(\mathbb{R}^3)} \leq C \langle \xi' \rangle^{m+2} \sigma^{2\mu m + \max(0, (m-1)/3)}, \quad j = 1, 2,$$

and

$$\langle \xi' \rangle \|\partial_{x_3} \phi_{*,k}\|_{H^m(\mathbb{R}^3)} + \|\theta \cdot \nabla_{x'} \partial_{x_3} \phi_{*,k}\|_{H^m(\mathbb{R}^3)} \leq C \langle \xi' \rangle^{m+1} \sigma^{2\mu(m+1) + \max(0, (m-1)/3)}.$$

Thus, we have $N_{0,\theta}(\partial_{x_j} \phi_{*,k}) \leq C \langle \xi' \rangle^2 \sigma^{2\mu}$ and $N_{2,\theta}(\partial_{x_j} \phi_{*,k}) \leq C \langle \xi' \rangle^4 \sigma^{6\mu+1/3}$, for $j = 1, 2, 3$, and consequently

$$\mathcal{N}_{\theta,\sigma}(\partial_{x_j} \phi_{*,k}) \leq C \langle \xi' \rangle^4 \sigma^{6\mu+1/3}, \quad j = 1, 2, 3.$$

according to (3.25). From this and (4.21), it then follows that

$$\mathcal{N}_{\theta,\sigma}(\phi_{*,k})\mathcal{N}_{\theta,\sigma}(\partial_{x_j}\phi_{*,k}) \leq C \langle \xi' \rangle^7 \sigma^{10\mu+2/3}, \quad j = 1, 2, 3. \quad (4.22)$$

Having seen this, we turn now to estimating $\widehat{\tilde{\rho}}_j$, where $\tilde{\rho}_j$ is defined by (4.1). As $A_\sigma^\sharp \in W^{\infty,\infty}(\mathbb{R}^3, \mathbb{R})^2$, we infer from (4.19) that $\phi_{*,k} \in \mathcal{C}_0^\infty(\mathbb{R}^3)$, and from (4.20) that $\text{supp } \phi_{*,k} \subset \mathcal{D}_R \times \mathbb{R}$. Thus, by performing the change of variable $y' = x' + t\theta \in \theta^\perp \oplus \mathbb{R}\theta$, in the following integral, we deduce from (4.18)-(4.19) that

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}^2} \phi_{*,k}^2(y', y_3) \mathcal{P}(\tilde{\rho}_j)(\theta, y', y_3) \exp \left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(y' + s\theta, y_3) ds \right) dy' dy_3 \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\theta^\perp} \phi_{*,k}^2(x' + t\theta, y_3) \mathcal{P}(\tilde{\rho}_j)(\theta, x' + t\theta, y_3) \exp \left(-i \int_{\mathbb{R}} \theta \cdot A_\sigma^\sharp(x' + s\theta, y_3) ds \right) dx' dt dy_3 \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\theta^\perp} h^2(t + r_{x'_k}) e^{-ix' \cdot \xi'} \varphi_k(x') \alpha_\sigma^2(y_3) \mathcal{P}(\tilde{\rho}_j)(\theta, x', y_3) dx' dt dy_3 \\ &= \int_{\mathbb{R}} \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \alpha_\sigma^2(y_3) \mathcal{P}(\tilde{\rho}_j)(\theta, y', y_3) dy' dy_3. \end{aligned} \quad (4.23)$$

Thus, taking $\mu > 0$ so small that $\kappa := 1/6 - 10\mu > 0$, we deduce from this, (4.4), and (4.22), that

$$\begin{aligned} & \left| \int_{\mathbb{R}} \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \alpha_\sigma^2(y_3) \mathcal{P}(\tilde{\rho}_j)(\theta, y', y_3) dy' dy_3 \right| \\ & \leq C \langle \xi' \rangle^7 \left(\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-\kappa} \right), \quad x_3 \in \mathbb{R}. \end{aligned} \quad (4.24)$$

Moreover, we see from (4.1) that $\tilde{\rho}_j \in \mathcal{C}^{0,1}(\mathbb{R}^3)$. Since $\text{supp } \tilde{\rho}_j \subset B(0, R) \times \mathbb{R}$, by Lemma 3.1, then $x \mapsto \mathcal{P}(\tilde{\rho}_j)(\theta, x) \in \mathcal{C}^{0,1}(\mathbb{R}^3)$, and we deduce from (3.18) upon making the substitution $s = \sigma^{2\mu}(x_3 - y_3)$ in the following integral, that

$$\begin{aligned} & \left| \int_{\mathbb{R}} \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \mathcal{P}(\tilde{\rho}_j)(\theta, y', y_3) \alpha_\sigma^2(y_3) dy' dy_3 - \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy' \right| \\ &= \left| \int_{\mathbb{R}} \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \alpha^2(s) \left(\mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3 - \sigma^{-2\mu}s) - \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) \right) dy' ds \right| \\ &\leq \int_{-1}^1 \int_{\theta^\perp \cap B(0, R+1)} \alpha^2(s) \left| \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3 - \sigma^{-2\mu}s) - \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) \right| dy' ds \leq C \sigma^{-2\mu}, \end{aligned}$$

for some constant $C > 0$ depending only on ω and M . Here, we used the fact that $\phi_{*,k}$ and α are supported in $B(0, R+1)$ and $(-1, 1)$, respectively. This and (4.24) yield

$$\left| \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy' \right| \leq C \langle \xi' \rangle^7 \left(\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-2\mu} + \sigma^{-\kappa} \right), \quad (4.25)$$

for all $x_3 \in \mathbb{R}$ and $k = 1, \dots, N$. Further, as A^\sharp is supported in $B(0, R) \times \mathbb{R}$ by assumption, it holds true that $\partial_{x_j} A^\sharp(y' + s\theta, x_3) = 0$ for all $s \in \mathbb{R}$, $x_3 \in \mathbb{R}$, and all $y' \in \theta^\perp$ such that $|y'| \geq R$. Therefore, we have

$$\mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) = 0, \quad y' \in \theta^\perp \cap (\mathbb{R}^2 \setminus B(0, R)), \quad x_3 \in \mathbb{R},$$

in virtue of (4.1), and hence

$$\int_{\theta^\perp} e^{-iy' \cdot \xi'} \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy' = \int_{\theta^\perp \cap B(0, R)} e^{-iy' \cdot \xi'} \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy', \quad x_3 \in \mathbb{R}.$$

In light of (4.12) and (4.16), this entails that

$$\widehat{\rho}_j(\xi', x_3) = \frac{1}{2\pi} \sum_{k=1}^N \int_{\theta^\perp \cap B(0, R)} e^{-iy' \cdot \xi'} \varphi_k(y') \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy', \quad x_3 \in \mathbb{R}. \quad (4.26)$$

Taking $\mu \in (0, 1/72]$, in such a way that we have $\kappa \geq 2\mu$, we infer from (4.25)-(4.26) that

$$\begin{aligned} \|\widehat{\rho}_j(\xi', \cdot)\|_{L^\infty(\mathbb{R})} &\leq \sum_{k=1}^N \left(\sup_{x_3 \in \mathbb{R}} \left| \int_{\theta^\perp} e^{-iy' \cdot \xi'} \varphi_k(y') \mathcal{P}(\tilde{\rho}_j)(\theta, y', x_3) dy' \right| \right) \\ &\leq C \langle \xi' \rangle^7 (\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-2\mu}), \quad x_3 \in \mathbb{R}. \end{aligned} \quad (4.27)$$

The last step of the proof is to notice from (4.1), (4.11), and the identity $\sum_{m=1,2} \theta_m \xi_m = \theta \cdot \xi' = 0$, that

$$\widehat{\rho}_j(\xi', x_3) = i \sum_{m=1,2} \theta_m \xi_j \widehat{a}_m(\xi', x_3) = i \sum_{m=1,2} \theta_m (\xi_j \widehat{a}_m - \xi_m \widehat{a}_j)(\xi', x_3), \quad x_3 \in \mathbb{R}, \quad j = 1, 2.$$

Thus, assuming that $\xi' = (\xi_1, \xi_2) \in \mathbb{R}^2 \setminus \{0\}$, we get from (4.13) upon choosing $\theta = (\xi_2/|\xi'|, -\xi_1/|\xi'|)$, that

$$\widehat{\rho}_j(\xi', x_3) = -\frac{\xi_j}{|\xi'|} \widehat{\beta}(\xi', x_3), \quad x_3 \in \mathbb{R}.$$

From this and (4.27), it then follows that

$$\|\widehat{\beta}(\xi', \cdot)\|_{L^\infty(\mathbb{R})} \leq \frac{|\xi_1| + |\xi_2|}{|\xi'|} \|\widehat{\beta}(\xi', \cdot)\|_{L^\infty(\mathbb{R})} \leq C \langle \xi' \rangle^7 (\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-2\mu}),$$

which yields (4.14) for $\xi' \neq 0$. Since $\widehat{\beta}(0, x_3) = 0$ for every $x_3 \in \mathbb{R}$, from (4.13), then (4.14) holds for $\xi' = 0$ as well, and the proof is complete. \square

Armed with Lemma 4.2, we turn now to proving the three main results of this paper.

5. PROOF OF THEOREMS 1.2, 1.3 AND 1.4

Let us start by reducing the analysis of the inverse problem under investigation to the case of transverse magnetic potentials. To do that, we consider $A' = (a'_i)_{1 \leq i \leq 3} \in \mathcal{A}$, and put $A := (a_1, a_2, 0)$, where

$$a_i(x', x_3) := a'_i(x', x_3) - \int_{-\infty}^{x_3} \partial_{x_i} a'_3(x', s) ds, \quad x = (x', x_3) \in \omega \times \mathbb{R}, \quad i = 1, 2. \quad (5.1)$$

Since $a'_3 \in C^3(\overline{\Omega})$ fulfills (1.9)-(1.10), from the very definition of \mathcal{A} , we have $a'_3 \in L^1_{x_3}(\mathbb{R}, H^3_0(\omega))$, where $H^3_0(\omega)$ denotes the closure of $\mathcal{C}^\infty_0(\omega)$ in $H^3(\omega)$. Thus, $e(x) := \int_{-\infty}^{x_3} a'_3(x', s) ds$ lies in $W^{3,\infty}(\Omega) \cap L^\infty_{x_3}(\mathbb{R}, H^3_0(\omega))$, and we deduce from the identity $A = A' - \nabla e$, arising from (5.1), that

$$dA' = dA, \text{ and } \Lambda_{A_* + A', q} = \Lambda_{A_* + A, q},$$

for all $A_* \in W^{2,\infty}(\Omega)^3$ and all $q \in W^{1,\infty}(\Omega)$. Moreover, it is easy to see that A obeys (1.9), in the sense that we have

$$\partial_x^\alpha A(x) = 0, \quad x \in \partial\Omega, \quad \alpha \in \mathbb{N}^3, \quad |\alpha| \leq 1. \quad (5.2)$$

Therefore, for each $A_* \in W^{2,\infty}(\Omega, \mathbb{R})^3$ and any $A_j \in A_* + \mathcal{A}$, $j = 1, 2$, we may assume without loss of generality, that the difference $A_2 - A_1$ reads

$$A = (a_1, a_2, 0), \quad (5.3)$$

and fulfills (5.2). We shall systematically do that in the sequel. For further reference, we put $A^\sharp := (a_1, a_2)$, where a_j , $j = 1, 2$, are extended by zero outside Ω . Summing up, we have

5.1. Proof of Theorem 1.2. We establish the uniqueness result $(dA_1, q_1) = (dA_2, q_2)$ in Subsection 5.1.1, while the proof of the stability estimate (1.13) can be found in Subsection 5.1.2.

5.1.1. Uniqueness result. For $\xi' = (\xi_1, \xi_2) \in \mathbb{R}^2 \setminus \{0\}$, we set $\xi'_\perp := (-|\xi'|^{-1}\xi_2, |\xi'|^{-1}\xi_1)$, and we decompose A^\sharp into the sum $(A^\sharp \cdot \xi')|\xi'|^{-2}\xi' + (A^\sharp \cdot \xi'_\perp)\xi'_\perp$, in such a way that the partial Fourier transform of $\partial_{x_3}A^\sharp$, reads

$$\partial_{x_3}\widehat{A^\sharp}(\xi', x_3) = \left(\frac{1}{2\pi} \int_{\mathbb{R}^2} e^{-ix' \cdot \xi'} \partial_{x_3}A^\sharp(x', x_3) \cdot \xi' dx' \right) \frac{\xi'}{|\xi'|^2} + i \frac{\partial_{x_3}\widehat{\beta}(\xi', x_3)}{|\xi'|} \xi'_\perp, \quad x_3 \in \mathbb{R}. \quad (5.4)$$

Next, recalling the hypothesis $\Lambda_{A_1, q_1} = \Lambda_{A_2, q_2}$, we get

$$\beta = \partial_{x_1}a_2 - \partial_{x_2}a_1 = 0 \text{ in } \Omega, \quad (5.5)$$

upon sending σ to infinity in (4.14). Moreover, we have $\nabla_{x'} \cdot \partial_{x_3}A^\sharp = \nabla \cdot \partial_{x_3}A = 0$, in virtue of (1.11), whence

$$\begin{aligned} \int_{\mathbb{R}^2} e^{-ix' \cdot \xi'} \partial_{x_3}A^\sharp(x', x_3) \cdot \xi' dx' &= i \int_{\mathbb{R}^2} \nabla_{x'} e^{-ix' \cdot \xi'} \cdot \partial_{x_3}A^\sharp(x', x_3) \\ &= -i \int_{\mathbb{R}^2} e^{-ix' \cdot \xi'} \nabla_{x'} \cdot \partial_{x_3}A^\sharp(x', x_3) dx' = 0. \end{aligned} \quad (5.6)$$

Putting this together with (5.4)-(5.5), we find that $|\xi'| \partial_{x_3}\widehat{A^\sharp}(\xi', x_3) = 0$ for a.e. $x_3 \in \mathbb{R}$. Since ξ' is arbitrary in $\mathbb{R}^2 \setminus \{0\}$, this entails that $\partial_{x_3}A^\sharp = 0$, and hence that $\partial_{x_3}a_1 = \partial_{x_3}a_2 = 0$ in \mathbb{R}^2 . From this, (5.5), and the fact that a_3 is uniformly zero, it then follows that $dA_1 = dA_2$.

Further, taking into account that $\partial_x^\alpha A_1 = \partial_x^\alpha A_2 = \partial_x^\alpha A_*$ on $\partial\Omega$, for every $\alpha \in \mathbb{N}^3$ such that $|\alpha| \leq 1$, we infer that $A \in W^{2,\infty}(\mathbb{R}^3, \mathbb{R})^3$. This and the identity $dA = 0$, yield $A = \nabla\Psi$, where the function $\Psi(x) = \int_0^1 x \cdot A(tx) dt$ lies in $W^{3,\infty}(\mathbb{R}^3, \mathbb{R})$. Moreover, since A vanishes in $\mathbb{R}^3 \setminus \Omega$, we may assume upon possibly adding a suitable constant, that the same is true for Ψ . Therefore, $\Psi|_{\partial\Omega} = 0$, and we find $\Lambda_{A_2, q_2} = \Lambda_{A_2 + \nabla\Psi, q_2} = \Lambda_{A_1, q_2}$, by combining the identity $A_1 = A_2 + \nabla\Psi$ with the gauge invariance property of the DN map. From this and the assumption $\Lambda_{A_1, q_1} = \Lambda_{A_2, q_2}$, it then follows that

$$\Lambda_{A_1, q_2} = \Lambda_{A_1, q_1}. \quad (5.7)$$

It remains to show that the function $q = q_2 - q_1$, duly extended by zero outside Ω , is uniformly zero in \mathbb{R}^3 . This can be done upon applying the orthogonality identity (3.50) with $A_1 = A_2$, i.e with $A = 0$ and $V = q$. In light of (5.7), we obtain that

$$\langle qu_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} = 0, \quad \sigma > \sigma_*. \quad (5.8)$$

Here $u_{j,\sigma}$, for $j = 1, 2$, is given by (3.3), and we have $(\overline{b_{1,\sigma}b_{2,\sigma}})(t, x) = 1$ for all $(t, x) \in (0, T) \times \mathbb{R}^3$, from (3.4), since $A_1 = A_2$.

Next, pick $\phi \in C_0^\infty(\mathbb{R}^3)$, with support in $\{x \in \mathbb{R}^3; |x| < 1\}$, and such that $\|\phi\|_{L^2(\mathbb{R}^3)}^2 = 1$. We fix $y \in \mathcal{D}_R(\theta) \times \mathbb{R}$, and choose $\delta > 0$ so small that $\phi_1(x) = \phi_2(x) := \delta^{-3/2}\phi(\delta^{-1}(x - y))$ is supported in $\mathcal{D}_R \times \mathbb{R}$. Thus, upon multiplying (5.8) by σ , and then sending σ to infinity, we find with the aid of (3.19) and (3.27), that

$$\int_0^{+\infty} \left(\int_{\mathbb{R}^3} q(\delta x' + y' + s\theta, \delta x_3 + y_3) |\phi(x', x_3)|^2 dx' dx_3 \right) ds = 0, \quad \delta > 0. \quad (5.9)$$

Actually, if $y' \in \mathcal{D}_R^-(\theta)$, then we have $|y' + s\theta| > R$ for any $s \leq 0$, and hence $q(\delta x' + y' + s\theta, \delta x_3 + y_3) = 0$, uniformly in $|x| < 1$, provided $\delta \in (0, 1)$. This and (5.9) yield that

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}^3} q(\delta x' + y' + s\theta, \delta x_3 + y_3) |\phi(x', x_3)|^2 dx' dx_3 \right) ds = 0, \quad \delta \in (0, 1), \quad (y', y_3) \in \mathcal{D}_R^-(\theta) \times \mathbb{R}. \quad (5.10)$$

By performing the change of variable $t = -s$ in the above integral, and then substituting $(-\theta)$ for θ in the resulting identity, we get that

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}^3} q(\delta x' + y' + s\theta, \delta x_3 + y_3) |\phi(x', x_3)|^2 dx' dx_3 \right) ds = 0, \quad \delta \in (0, 1), \quad (y', y_3) \in \mathcal{D}_R^-(\theta) \times \mathbb{R}.$$

This and (5.10) yield that

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}^3} q(\delta x' + y' + s\theta, \delta x_3 + y_3) |\phi(x', x_3)|^2 dx' dx_3 \right) ds = 0, \quad \delta \in (0, 1), \quad (y', y_3) \in \mathcal{D}_R \times \mathbb{R}.$$

Next, sending δ to zero in the above identity, and taking into account that ϕ is normalized in $L^2(\mathbb{R}^3)$, we obtain for each $\theta \in \mathbb{S}^1$, that

$$\mathcal{P}(q)(\theta, y', y_3) = \int_{\mathbb{R}} q(y' + s\theta, y_3) ds = 0, \quad (y', y_3) \in \mathcal{D}_R \times \mathbb{R}.$$

This entails $q = 0$, since the partial X-ray transform is injective.

5.1.2. Proof of the stability estimate (1.13). We have

$$|\xi'| \left| \partial_{x_3} \widehat{A}^\sharp(\xi', x_3) \right| = \left| \partial_{x_3} \widehat{\beta}(\xi', x_3) \right|, \quad \xi' \in \mathbb{R}^2, \quad x_3 \in \mathbb{R},$$

by (5.4) and (5.6), so we infer from (4.14)-(4.15) that

$$|\widehat{\beta}(\xi', x_3)| + |\xi'| |\partial_{x_3} \widehat{A}^\sharp(\xi', x_3)| \leq C \langle \xi' \rangle^8 (\sigma^6 \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{-\epsilon}), \quad \xi' \in \mathbb{R}^2, \quad x_3 \in \mathbb{R}, \quad (5.11)$$

for all $\sigma > \sigma_*$, the constants C and ϵ being the same as in Lemma 4.2.

Fix $\rho \in (1, +\infty)$ and put $\mathcal{C}_\rho := \{\xi' \in \mathbb{R}^2; \rho^{-1} \leq |\xi'| \leq \rho\}$. Then, upon applying the Plancherel theorem, we obtain

$$\begin{aligned} \|\partial_{x_3} a_j(\cdot, x_3)\|_{L^2(\omega)}^2 &\leq \|\partial_{x_3} \widehat{a}_j(\xi', x_3)\|_{L^2(B(0, \rho^{-1}))}^2 + \rho^{-2} \int_{\mathbb{R}^2 \setminus B(0, \rho)} \langle \xi' \rangle^2 |\partial_{x_3} \widehat{a}_j(\xi', x_3)|^2 d\xi' \\ &\quad + \|\partial_{x_3} \widehat{a}_j(\xi', x_3)\|_{L^2(\mathcal{C}_\rho)}^2, \quad x_3 \in \mathbb{R}, \quad j = 1, 2. \end{aligned} \quad (5.12)$$

Further, as $\|\partial_{x_3} \widehat{a}_j(\xi', x_3)\|_{L^2(B(0, \rho^{-1}))}^2 \leq |\omega| \rho^{-2} \|a_j\|_{W^{1, \infty}(\Omega)}^2$ and $\int_{\mathbb{R}^2 \setminus B(0, \rho)} \langle \xi' \rangle^2 |\partial_{x_3} \widehat{a}_j(\xi', x_3)|^2 d\xi' \leq \|a_j\|_{W^{1, \infty}(\Omega)}^2$, then there exists a constant $C > 0$, depending only on M and ω , such that we have

$$\|\partial_{x_3} \widehat{a}_j(\xi', x_3)\|_{L^2(B(0, \rho^{-1}))}^2 + \rho^{-2} \int_{\mathbb{R}^2 \setminus B(0, \rho)} \langle \xi' \rangle^2 |\partial_{x_3} \widehat{a}_j(\xi', x_3)|^2 d\xi' \leq \frac{M}{\rho^2}, \quad (5.13)$$

according to (1.12). On the other hand, we derive from (5.11) that

$$|\partial_{x_3} \widehat{a}_j(\xi', x_3)|^2 \leq C \rho^{14} (\sigma^{12} \delta^2 + \sigma^{-2\epsilon}), \quad \xi' \in \mathcal{C}_\rho \cap B(0, \rho), \quad x_3 \in \mathbb{R}, \quad \sigma > \sigma_*, \quad j = 1, 2,$$

where $\delta := \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|$. Putting this and (5.12)-(5.13) together, we get for every $\sigma > \sigma_*$, that

$$\|\partial_{x_3} a_j\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))}^2 \leq C (\rho^{16} \sigma^{12} \delta^2 + \rho^{16} \sigma^{-2\epsilon} + \rho^{-2}), \quad x_3 \in \mathbb{R}, \quad j = 1, 2. \quad (5.14)$$

Now, choosing ρ so large that $\rho > \sigma_*^{\epsilon/8}$, we get upon taking $\sigma = \rho^{8/\epsilon} > \sigma_*$ in (5.14), that

$$\|\partial_{x_3} a_j(\cdot, x_3)\|_{L^2(\omega)}^2 \leq C (\rho^{M_\epsilon} \delta^2 + \rho^{-2}), \quad x_3 \in \mathbb{R}, \quad j = 1, 2, \quad (5.15)$$

where $M_\epsilon := 16 + 96/\epsilon$. Thus, if $\delta < \delta_0 := \sigma_*^{-\epsilon(M_\epsilon+2)/16}$, then we have $\delta^{-2/(M_\epsilon+2)} > \sigma_*^{\epsilon/8}$, and we may apply (5.15) with $\rho = \delta^{-2/(M_\epsilon+2)}$, getting

$$\|\partial_{x_3} a_j\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))}^2 \leq 2C\delta^{2\mu_0}, \text{ with } \mu_0 := \frac{2}{M_\epsilon + 2} \in (0, 1), \quad j = 1, 2. \quad (5.16)$$

From this and the fact, arising from (1.12), that $\|\partial_{x_3} a_j\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} \leq (2M\delta_0^{-2\mu_0})\delta^{2\mu_0}$ for all $\delta \geq \delta_0$, it then follows that (5.16) remains valid for every $\delta > 0$.

Finally, arguing as before with β instead of $\partial_{x_3} A^\sharp$, we obtain in a similar way from (5.11), that $\|\beta\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))}$ is upper bounded, up to some multiplicative constant depending only on M and ω , by δ^{μ_0} , and hence (1.13) follows from this and (5.16).

5.2. Proof of Theorem 1.3. The proof is an adaptation of the one of (1.2), where the adaptation is to take into account the extra information given by (1.14). Actually, since $A_j = (a_{1,j}, a_{2,j}, a_{3,*})$ and $A = (A^\sharp, 0)$ with $A^\sharp = (a_1, a_2)$, by (5.3), then (1.14) yields

$$\|\partial_{x_1} a_2 - \partial_{x_2} a_1\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} = \|\partial_{x_1} a_2 - \partial_{x_2} a_1\|_{L_{x_3}^\infty(-r, r; L^2(\omega))}, \quad (5.17)$$

and

$$\|\partial_{x_3} a_j\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))} = \|\partial_{x_3} a_j\|_{L_{x_3}^\infty(-r, r; L^2(\omega))}, \quad j = 1, 2. \quad (5.18)$$

More precisely, we still consider GO solutions $u_{1,\sigma}$ and $u_{2,\sigma}$, defined by (3.3)-(3.4) and (3.22), with $\phi_1 = \partial_{x_j} \phi$, for $j = 1, 2, 3$, and $\phi_2 = \phi$, where ϕ is given by (4.18)-(4.19). The parameter x_3 appearing in (4.18), is taken in $(-r, r)$, and we impose $\sigma > (r' - r)^{-24}$, in such a way that $\phi \in C_0^\infty(\mathcal{D}_R^-(\theta) \times (-r', r'))$. Moreover, the functions

$$f_\sigma(t, x) = \Phi_2(2\sigma t, x) b_2(2\sigma t, x) e^{i\sigma(x\theta - \sigma t)} \text{ and } g_\sigma = \Phi_1(2\sigma t, x) b_1(2\sigma t, x) e^{i\sigma(x\theta - \sigma t)}, \quad (t, x) \in \Sigma,$$

lie in $H_0^{2,1}((0, T) \times \Gamma_{r'})$, and we infer from (3.50) upon arguing as the derivation of Lemma 4.2, that

$$\|\widehat{\beta}(\xi', \cdot)\|_{L^\infty(-r', r')} \leq C \langle \xi' \rangle^7 (\sigma^6 \|\Lambda_{A_1, q_1, r'} - \Lambda_{A_2, q_2, r'}\| + \sigma^{-\epsilon}),$$

and that

$$\|\partial_{x_3} \widehat{\beta}(\xi', \cdot)\|_{L^\infty(r', r')} \leq C \langle \xi' \rangle^8 (\sigma^6 \|\Lambda_{A_1, q_1, r'} - \Lambda_{A_2, q_2, r'}\| + \sigma^{-\epsilon}).$$

for all $\xi' \in \mathbb{R}^2$ and some $\epsilon > 0$. Here, the constant C depends only on ω , T , M , r , r' and ϵ . The desired result follows from this and (5.17)-(5.18) by arguing in the same way as in the proof of Theorem 1.2.

5.3. Proof of Theorem 1.4. We only prove (1.16), the derivation of (1.19) being quite similar to the one of (1.15). To this end, we fix $\xi' \in \mathbb{R}^2$, and remind that $A = (A^\sharp, 0) \in \mathcal{A}_0$, with $A^\sharp = (a_1, a_2)$, satisfies $\partial_{x_1} a_1 + \partial_{x_2} a_2 = 0$ in \mathbb{R}^2 , so we get

$$\begin{aligned} \widehat{A^\sharp}(\xi', x_3) \cdot \xi' &= i(2\pi)^{-1} \int_{\mathbb{R}^2} A^\sharp(x', x_3) \cdot \nabla_{x'} e^{-ix' \cdot \xi'} dx' \\ &= -i(2\pi)^{-1} \int_{\mathbb{R}^2} e^{-ix' \cdot \xi'} (\partial_{x_1} a_1 + \partial_{x_2} a_2)(x', x_3) dx' = 0, \quad x_3 \in \mathbb{R}, \end{aligned}$$

upon integrating by parts. Thus, remembering that $\xi'_\perp = (-|\xi'|^{-1}\xi_2, |\xi'|^{-1}\xi_1)$ whenever $\xi' \neq 0$, we obtain

$$\widehat{A^\sharp}(\xi', x_3) = (\widehat{A^\sharp}(\xi', x_3) \cdot \xi'_\perp) \xi'_\perp = (-\xi_2 \widehat{a_1} + \xi_1 \widehat{a_2})(\xi', x_3) \frac{\xi'_\perp}{|\xi'|}, \quad x_3 \in \mathbb{R},$$

and consequently

$$|\xi'| \widehat{A^\sharp}(\xi', x_3) = -i \widehat{\beta}(\xi', x_3), \quad x_3 \in \mathbb{R}, \quad (5.19)$$

from (4.13), the above identity being valid for $\xi' = 0$ as well. Therefore, arguing as in the derivation of (1.13) from (4.14), we infer from (4.15) and (5.19) that

$$\|A\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega))}^3 \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{\mu_1}, \quad (5.20)$$

where $C > 0$ and $\mu_1 \in (0, 1)$ are two constants depending only on T , ω , and M .

We turn now to estimating $\|q\|_{L_{x_3}^\infty(\mathbb{R}, H^{-1}(\omega))^3}$, where $q = q_1 - q_2$. With reference to (4.16)-(4.17), we fix $x_3 \in \mathbb{R}$ and $k \in \{1, \dots, N\}$, and pick a function $\phi_{*,k}$, expressed by (4.18)-(4.19) in the particular case where A_σ^\sharp is uniformly zero, i.e.

$$\phi_{*,k}(y) := h\left(y' \cdot \theta + r_{x'_k}\right) e^{-\frac{i}{2}y' \cdot \xi'} \varphi_k^{1/2}(y' - (y' \cdot \theta)\theta) \alpha_\sigma(y_3), \quad y' \in \mathbb{R}^2, \quad y_3 \in \mathbb{R}. \quad (5.21)$$

In view of Proposition 3.1, we consider a GO solution $u_{j,\sigma}$, $j = 1, 2$, to the magnetic Schrödinger equation $(i\partial_t + \Delta_{A_j} + q_j)u_{j,\sigma} = 0$ in Q , described by (3.3) with

$$\Phi_1 = \overline{\Phi_{*,k}}, \quad \Phi_2 = \Phi_{*,k}, \quad \text{and} \quad \Phi_{*,k}(t, x) := \phi_{*,k}(x' - t\theta, x_3), \quad t \in \mathbb{R}, \quad x' \in \mathbb{R}^2, \quad x_3 \in \mathbb{R}. \quad (5.22)$$

Bearing in mind that $\nabla \cdot A = 0$, we then apply (3.50) with $V = q - A \cdot (A_1 + A_2)$, getting

$$\langle qu_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} = \langle A \cdot ((A_1 + A_2)u_{2,\sigma} - 2i\nabla u_{2,\sigma}), u_{1,\sigma} \rangle_{L^2(Q)} + \langle (\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2})f_\sigma, g_\sigma \rangle_{L^2(\Sigma)},$$

where f_σ and g_σ are given by (3.47) and (3.51), respectively. Thus, we have

$$|\langle qu_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)}| \leq C \|A\|_{L^\infty(\Omega)^3} \|u_{1,\sigma}\|_{L^2(Q)} \|u_{2,\sigma}\|_{L^2(0,T;H^1(\Omega))} + |\langle (\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2})f_\sigma, g_\sigma \rangle_{L^2(\Sigma)}|,$$

and consequently

$$|\langle qu_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)}| \leq C \sigma^{8\mu} \left(\|A\|_{L^\infty(\Omega)^3} + \sigma^{17/3} \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| \right) \langle \xi' \rangle^6, \quad (5.23)$$

by (3.38), (3.55), and (4.21).

On the other hand, it follows readily from (3.3) and (5.22), that

$$\langle qu_{2,\sigma}, u_{1,\sigma} \rangle_{L^2(Q)} = \int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) (\overline{b_{1,\sigma} b_{2,\sigma}})(2\sigma t, y) dy dt + R_{k,\sigma}, \quad (5.24)$$

where $b_{j,\sigma}$, $j = 1, 2$, is given by (3.4), and

$$\begin{aligned} R_{k,\sigma} &:= \int_Q q(y) \Phi_{*,k}(2\sigma t, y) \left(b_{2,\sigma}(2\sigma t, y) e^{i\sigma(y' \cdot \theta - \sigma t)} \overline{\psi_{1,\sigma}}(t, y) + \psi_{2,\sigma}(t, y) \overline{b_{1,\sigma}}(2\sigma t, y) e^{-i\sigma(y' \cdot \theta - \sigma t)} \right) dy dt \\ &\quad + \int_Q q(y) (\psi_{2,\sigma} \overline{\psi_{1,\sigma}})(t, y) dy dt. \end{aligned}$$

Therefore, $|R_{k,\sigma}|$ is majorized by

$$\begin{aligned} &\|q\|_{L^\infty(\Omega)} \left(\|\Phi_{*,k}(2\sigma \cdot, \cdot)\|_{L^2(Q)} (\|\psi_{1,\sigma}\|_{L^2(Q)} + \|\psi_{2,\sigma}\|_{L^2(Q)}) + \|\psi_{1,\sigma}\|_{L^2(Q)} \|\psi_{2,\sigma}\|_{L^2(Q)} \right) \\ &\leq C \sigma^{-11/6} \mathcal{N}_{\theta,\sigma}(\phi_{*,k})^2, \end{aligned}$$

in virtue of (3.25)–(3.27) and (3.39), so we infer from (4.21) that

$$|R_{k,\sigma}| \leq C \sigma^{8\mu-7/6} \langle \xi' \rangle^6. \quad (5.25)$$

We turn now to examining the first term in the right hand side of (5.24). In light of (3.4), we have

$$\int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) (\overline{b_{1,\sigma} b_{2,\sigma}})(2\sigma t, y) dy dt = \int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) dy dt + r_{k,\sigma}, \quad (5.26)$$

with

$$r_{k,\sigma} := \int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) \left(e^{-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(y' - s\theta, y_3) ds} - 1 \right) dy dt. \quad (5.27)$$

Next, as $e^{-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(y' - s\theta, y_3) ds} - 1 = -i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(y' - \tau\theta, y_3) e^{-i \int_0^\tau \theta \cdot A_\sigma^\sharp(y' - s\theta, y_3) ds} d\tau$, we have

$$\left| e^{-i \int_0^{2\sigma t} \theta \cdot A_\sigma^\sharp(y' - s\theta, y_3) ds} - 1 \right| \leq 2\sigma T \|A_\sigma^\sharp\|_{L^\infty(\mathbb{R}^3)^2} \leq C\sigma \|A\|_{L^\infty(\Omega)^3}, \quad (t, y) \in Q.$$

Here we used the fact, arising from (3.9) and (3.15)-(3.16), that for any $\sigma > 0$, $\|A_\sigma^\sharp\|_{L^\infty(\mathbb{R}^3)^2}$ is majorized, up to some multiplicative constant that is independent of σ , by $\|A^\sharp\|_{L^\infty(\Omega)^2}$. Therefore, we infer from (1.12), (3.39), and (4.21), that

$$|r_{k,\sigma}| \leq C\sigma \|A\|_{L^\infty(\mathbb{R}^3)^3} \|\Phi_{*,k}(2\sigma \cdot, \cdot)\|_{L^2(Q)}^2 \leq C\sigma \|A\|_{L^\infty(\mathbb{R}^3)^3} \|\phi_{*,k}\|_{L^2(\mathbb{R}^3)}^2 \leq C\sigma \|A\|_{L^\infty(\mathbb{R}^3)^3} \langle \xi' \rangle^6 \sigma^{8\mu}. \quad (5.28)$$

We are left with the task of examining the integral

$$\begin{aligned} \int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) dy dt &= \int_0^T \int_{\mathbb{R}^3} q(y) \phi_{*,k}^2(y' - 2\sigma t\theta, y_3) dy' dy_3 dt \\ &= \frac{1}{2\sigma} \int_0^{2\sigma T} \int_{\mathbb{R}^3} q(y' + s\theta, y_3) \phi_{*,k}^2(y) dy' dy_3 ds, \end{aligned} \quad (5.29)$$

appearing in the right hand side of (5.26). To do that, we notice for all $\sigma > \sigma_*$, that

$$q(y' + s\theta, y_3) \phi_{*,k}^2(y) = 0, \quad s \in (-\infty, 0) \cup (2\sigma T, +\infty), \quad y' \in \mathbb{R}^2, \quad y_3 \in \mathbb{R},$$

since q and $\phi_{*,k}$ are supported in $B(0, R) \times \mathbb{R}$ and $\mathcal{D}_R^-(\theta) \times \mathbb{R}$, respectively, and that $|y' + s\theta| > R$ whenever $y' \in \mathcal{D}_R^-(\theta)$ and $s \in (-\infty, 0) \cup (2\sigma T, +\infty)$. In view of (4.3) and (5.29), this entails that

$$\begin{aligned} \int_Q q(y) \Phi_{*,k}^2(2\sigma t, y) dy dt &= \frac{1}{2\sigma} \int_{\mathbb{R}} \int_{\mathbb{R}^3} q(y' + s\theta, y_3) \phi_{*,k}^2(y) dy' dy_3 ds \\ &= \frac{1}{2\sigma} \int_{\mathbb{R}^3} \mathcal{P}(q)(\theta, y', y_3) \phi_{*,k}^2(y) dy' dy_3. \end{aligned}$$

Thus, arguing in the same way as in the derivation of (4.23), we infer from (5.21) that

$$|\widehat{q}(\xi', x_3)| \leq C \langle \xi' \rangle^6 \left(\sigma^{20/3} \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{8\mu+1} \|A\|_{L^\infty(\Omega)^3} + \sigma^{8\mu-1/6} \right), \quad (5.30)$$

for every $\sigma > \sigma_*$.

The next step of the proof is to upper bound $\|A\|_{L^\infty(\Omega)^3}$ in terms of $\|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|$. To do that, we pick $p > 2$ and apply Sobolev's embedding theorem (see e.g. [13, Corollary IX.14]), getting $\|A(\cdot, x_3)\|_{L^\infty(\omega)^3} \leq C \|A(\cdot, x_3)\|_{W^{1,p}(\omega)^3}$ for a.e. $x_3 \in \mathbb{R}$, where the constant $C > 0$ depends only on ω . Interpolating, we thus obtain that

$$\|A(\cdot, x_3)\|_{L^\infty(\omega)^3} \leq C \|A(\cdot, x_3)\|_{W^{2,p}(\omega)^3}^{1/2} \|A(\cdot, x_3)\|_{L^p(\omega)^3}^{1/2}, \quad x_3 \in \mathbb{R}.$$

This and (5.20) yield

$$\|A\|_{L^\infty(\Omega)^3} \leq C \|A\|_{L_{x_3}^\infty(\mathbb{R}, L^p(\omega)^3)}^{1/2} \leq C \|A\|_{L_{x_3}^\infty(\mathbb{R}, L^2(\omega)^3)}^{1/p} \leq C \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{\mu_1/p},$$

for some constant $C > 0$ depending only on ω , M and T . Then, we find by substituting the right hand side of the above estimate for $\|A\|_{L^\infty(\Omega)^3}$ in (5.30), that

$$|\widehat{q}(\xi', x_3)| \leq C \langle \xi' \rangle^6 \left(\sigma^{20/3} \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| + \sigma^{8\mu+1} \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\|^{\mu_1/p} + \sigma^{8\mu-1/6} \right), \quad \sigma > \sigma_*. \quad (5.31)$$

With the notations of Subsection 5.1.2, we infer from (5.31) and the estimate

$$\int_{\mathbb{R}^2 \setminus B(0, \rho)} \langle \xi' \rangle^{-2} |\widehat{q}(\xi', x_3)|^2 d\xi' \leq \frac{M}{\rho^2}, \quad x_3 \in \mathbb{R},$$

which holds true for any $\rho \in (1, +\infty)$, that

$$\|q\|_{L^\infty_{x_3}(\mathbb{R}, H^{-1}(\omega))} \leq C \left(\rho^6 \sigma^{20/3} \delta^{\mu_1/p} + \sigma^{8\mu-1/6} + \rho^{-1} \right), \quad \sigma > \sigma_*, \quad (5.32)$$

where $\delta = \|\Lambda_{A_1, q_1} - \Lambda_{A_2, q_2}\| \in (0, 1)$. Thus, for $\mu \in (0, 1/48)$ and $\delta \in \left(0, \sigma_*^{-(41-48\mu)p/(6\mu_1)}\right)$, we obtain (1.16) with $\mu_2 := (1 - 48\mu)\mu_1/(7p(41 - 48\mu))$ by taking $\rho = \delta^{-\mu_2}$ and $\sigma = \delta^{-42\mu_2/(1-48\mu)}$ in (5.32).

REFERENCES

- [1] S. Avdonin, S. Lenhart and V. Protopopescu, *Determining the potential in the Schrödinger equation from the Dirichlet to Neumann map by the Boundary Control method*, J. Inverse and Ill-Posed Problems, 13 (2005), no. 5, 317-330.
- [2] C. Bardos, G. Lebeau and J. Rauch, *Sharp sufficient conditions for the observation, control and stabilization from the boundary*, SIAM J. Control Optim. 30 (1992), 1024-1165.
- [3] L. Baudouin and J.-P. Puel, *Uniqueness and stability in an inverse problem for the Schrödinger equation*, Inverse Problems 18 (2002), 1537-1554.
- [4] M. Belishev, *An approach to multidimensional inverse problems for the wave equation*, Dokl. Akad. Nauk SSSR, 297 (1987), 524-527.
- [5] M. Bellassoued, *Uniqueness and stability in determining the speed of propagation of second-order hyperbolic equation with variable coefficients*, Appl. Anal. 83 (2004), 983-1014.
- [6] M. Bellassoued, *Stable determination of coefficients in the dynamical Schrödinger equation in a magnetic field*, preprint, arXiv:1510.04247.
- [7] M. Bellassoued and H. Benjoud, *Stability estimate for an inverse problem for the wave equation in a magnetic field*, Appl. Anal. 87 (2008), no. 3, 277-292.
- [8] M. Bellassoued and M. Choulli, *Logarithmic stability in the dynamical inverse problem for the Schrödinger equation by an arbitrary boundary observation*, J. Math. Pures Appl., 91 (2009), no. 3, 233-255.
- [9] M. Bellassoued and M. Choulli, *Stability estimate for an inverse problem for the magnetic Schrödinger equation from the Dirichlet-to-Neumann map*, J. Funct. Anal., 258 (2010), 161-195.
- [10] M. Bellassoued and D. Dos Santos Ferreira, *Stable determination of coefficients in the dynamical anisotropic Schrödinger equation from the Dirichlet-to-Neumann map*, Inverse Problems, 26 (2010), 125010.
- [11] M. Bellassoued, D. Jellali and M. Yamamoto, *Lipschitz stability for a hyperbolic inverse problem by finite local boundary data*, Applicable Analysis 85 (2006), 1219-1243.
- [12] M. Bellassoued, Y. Kian and E. Soccorsi, *An inverse stability result for non compactly supported potentials by one arbitrary lateral Neumann observation*, Journal of Differential Equations 260 (2016), no. 10, 7535-7562.
- [13] H. Brezis, *Analyse Fonctionnelle. Théorie et applications*, Collection mathématiques appliquées pour la maîtrise, Masson, Paris 1993.
- [14] A. L. Bukhgeim and M. V. Klibanov, *Global uniqueness of class of multidimensional inverse problems*, Soviet Math. Dokl., 24 (1981), 244-247.
- [15] F. Cardoso and R. Mendoza, *On the hyperbolic Dirichlet to Neumann functional*, Commun. Partial Diff. Equat., 21 (1996), 1235-1252.
- [16] T. Cazenave and A. Haraux, *Introduction aux problèmes d'évolution semi-linéaires*, ellipses, Paris, 1990.
- [17] M. Choulli, E. Soccorsi, *Recovering the twisting function in a twisted waveguide from the DN map*, J. Spectr. Theory, 5 (2015), no. 2, 295-329.
- [18] M. Choulli, Y. Kian and E. Soccorsi, *Determining the time dependent external potential from the DN map in a periodic quantum waveguide*, SIAM J. Math. Anal., 47 (2015), no. 6, 4536-4558.
- [19] M. Choulli, Y. Kian and E. Soccorsi, *Stability result for elliptic inverse periodic coefficient problem by partial Dirichlet-to-Neumann map*, arXiv:1601.05355.
- [20] M. Choulli, Y. Kian and E. Soccorsi, *On the Calderón problem in periodic cylindrical domain with partial Dirichlet and Neumann data*, arXiv:1601.05358.
- [21] P.-Y. Chang and H.-H. Lin, *Conductance through a single impurity in the metallic zigzag carbon nanotube*, Appl. Phys. Lett. 95, (2009), 082104.
- [22] M. Cristofol and E. Soccorsi, *Stability estimate in an inverse problem for non-autonomous magnetic Schrödinger equations*, Applicable Analysis, 90 (2011), no. 10, 1499-1520.
- [23] D. Dos Santos Ferreira, C. E. Kenig, J. Sjöstrand and G. Uhlmann, *Determining a magnetic Schrödinger operator from partial Cauchy data*, Comm. Math. Phys., 271 (2007), no. 2, 467-488.

- [24] G. Eskin, *Inverse problems for the Schrödinger equations with time-dependent electromagnetic potentials and the Aharonov-Bohm effect*, J. Math. Phys., 49 (2008), no. 2, 022105, 18 pp.
- [25] G. Eskin, *Global uniqueness in the inverse scattering problem for the Schrödinger operator with external Yang-Mills potentials*, Comm. Math. Phys., 222 (2001), no. 3, 503-531.
- [26] G. Eskin and J. Ralston, *Inverse scattering problem for the Schrödinger equation with magnetic potential at a fixed energy*, Commun. Math. Phys., 173 (1995), 199-224.
- [27] P. Grisvard, *Elliptic problems in nonsmooth domains*, Pitman, London, 1985.
- [28] V. Isakov and Z. Sun, *Stability estimates for hyperbolic inverse problems with local boundary data*, Inverse Problems 8 (1992), 193-206.
- [29] C. Kane, L. Balents and M. P. A. Fisher, *Coulomb Interactions and Mesoscopic Effects in Carbon Nanotubes*, Phys. Rev. Lett., 79, (1997), 5086-5089.
- [30] O. Kavian, Y. Kian and E. Soccorsi, *Uniqueness and stability results for an inverse spectral problem in a periodic waveguide*, Journal de Mathématiques Pures et Appliquées 104 (2015), no. 6, 1160-1189.
- [31] Y. Kian, *Stability of the determination of a coefficient for wave equations in an infinite waveguide*, Inverse Probl. Imaging, 8 (2014), no. 3, 713-732.
- [32] Y. Kian, *A multidimensional Borg-Levinson theorem for magnetic Schrödinger operators with partial spectral data*, arXiv:1504.04514.
- [33] Y. Kian, Q. S. Phan and E. Soccorsi, *Carleman estimate for infinite cylindrical quantum domains and application to inverse problems*, Inverse Problems, 30, 5 (2014), 055016.
- [34] Y. Kian, Q. S. Phan and E. Soccorsi, *Hölder stable determination of a quantum scalar potential in unbounded cylindrical domains*, Jour. Math. Anal. Appl., 426 (2015), no. 1, 194-210.
- [35] M. V. Klibanov, *Inverse problems and Carleman estimates*, Inverse Problems 8 (1992), 575-596.
- [36] M. V. Klibanov and A. A. Timonov, *Carleman estimates for coefficient inverse problems and numerical applications*, VSP, Utrecht, 2004.
- [37] K. Krupchyk, M. Lassas and G. Uhlmann, *Inverse problems with partial data for a magnetic Schrödinger operator in an infinite slab and on a bounded domain*, Comm. Math. Phys., 312 (2012), 87-126.
- [38] K. Krupchyk and G. Uhlmann, *Uniqueness in an inverse boundary problem for a magnetic Schrödinger operator with a bounded magnetic potential*, Comm. Math. Phys., 327 (2014), no. 1, 993-1009.
- [39] X. Li and G. Uhlmann, *Inverse problems with partial data in a slab*, Inverse Problems and Imaging, 4 (2010), no. 3, 449-462.
- [40] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications*, Vol. I, Dunod, Paris, 1968.
- [41] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications*, Vol. II, Dunod, Paris, 1968.
- [42] E. Machtyngier, *Exact controllability for the Schrödinger equation*, SIAM J. Control and Optimization, 32 (1994), no. 1, 24-34.
- [43] Rakesh, *An inverse problem for the wave equation in the half plane*, Inverse Problems, 9 (1993), 433-441.
- [44] Rakesh and W. Symes, *Uniqueness for an inverse problems for the wave equation*, Commun. Partial Diff. Equat., 13 (1988), 87-96.
- [45] M. Salo, *Inverse problems for nonsmooth first order perturbations of the Laplacian*, Ann. Acad. Scient. Fenn. Math. Dissertations, Vol. 139, 2004.
- [46] M. Salo, *Semiclassical pseudodifferential calculus and the reconstruction of a magnetic field*, Commun. Partial Diff. Equat., 31 (2006), no. 11, 1639-1666.
- [47] P. Stefanov and G. Uhlmann, *Stability estimates for the hyperbolic Dirichlet to Neumann map in anisotropic media*, J. Funct. Anal., 154 (1998), 330-358.
- [48] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, 1970.
- [49] Z. Sun, *On continuous dependence for an inverse initial boundary value problem for the wave equation*, J. Math. Anal. Appl., 150 (1990) 188-204.
- [50] Z. Sun, *An inverse boundary value problem for the Schrödinger operator with vector potentials*, Trans. Amer. Math. Soc., 338 (1992), no. 2, 953-969.
- [51] H. Tanabe, *Functional Analytic Methods for Partial Differential Equations*, Marcel Dekker, (1996).
- [52] H. Triebel, *Interpolation Theory Function Spaces Differential Operators*, Johann Ambrosius Barth Verlag Heidelberg-Leipzig, Germany, 1995.
- [53] L. Tzou, *Stability estimates for coefficients of magnetic Schrödinger equation from full and partial boundary measurements*, Commun. Partial Diff. Equat., 11 (2008), 1911-1952.
- [54] Y. Yang, *Determining the first order perturbation of a bi-harmonic operator on bounded and unbounded domains from partial data*, J. Differential Equations, 257 (2014), 3607-3639.

UNIVERSITY OF TUNIS EL MANAR, NATIONAL ENGINEERING SCHOOL OF TUNIS, ENIT-LAMSIN, B.P. 37, 1002 TUNIS, TUNISIA

E-mail address: mourad.bellassoued@fsb.rnu.tn

AIX-MARSEILLE UNIVERSITÉ, CNRS, CPT UMR 7332, 13288 MARSEILLE & UNIVERSITÉ DE TOULON, CNRS, CPT UMR 7332, 83957 LA GARDE, FRANCE.

E-mail address: yavar.kian@univ-amu.fr

AIX-MARSEILLE UNIVERSITÉ, CNRS, CPT UMR 7332, 13288 MARSEILLE & UNIVERSITÉ DE TOULON, CNRS, CPT UMR 7332, 83957 LA GARDE, FRANCE.

E-mail address: eric.soccorsi@univ-amu.fr